

MULTIJURISDICTIONAL SYPHILIS OUTBREAK SURVEILLANCE REPORT: JULY 2018

This surveillance report has been authorised by Health Departments in outbreak affected jurisdictions and the Office of Health Protection.

The Multijurisdictional Syphilis Outbreak (MJSO) Working Group was formed by the Communicable Diseases Network of Australia (CDNA) in April 2015, in response to an ongoing outbreak of infectious syphilis among Aboriginal and Torres Strait Islander people living largely in remote and rural areas of northern Australia. Increased notifications associated with the outbreak in northern Australia were first reported in January 2011 in northwest Queensland (Qld), followed by the Northern Territory (NT) in July 2013, and the Kimberley region of Western Australia (WA) in June 2014. In March 2017, South Australia (SA) declared an outbreak in the Far North and Western and Eyre regions from November 2016.

This surveillance report summarises the outbreak epidemiological data as of 30 June 2018 (current status).

Please note: the MJSO Surveillance Report was formerly known as the MJSO Communique. Historical epidemiological data and MJSO activities are published in earlier MJSO Communiques available at <http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-infectious-syphilis-outbreak.htm>

Current status

Outbreak data to 30 June 2018[‡] are summarised in Figure 1, Table 1 and Table 2 below.

Outbreak cases are reported as either category 1 or category 2: category 1 cases include Aboriginal and Torres Strait Islander people residing in an outbreak declared region[‡] at the time of diagnosis, and; category 2 cases include people who are a sexual contact of a confirmed outbreak case which includes Aboriginal and Torres Strait Islander people who do not reside in an outbreak area at the time of diagnosis and non-Indigenous people regardless of where they reside.

Between 1 – 30 June 2018[‡], there were 54 outbreak cases (category 1 and 2 cases) reported from the four outbreak affected jurisdictions: 22 in Queensland; 26 in the Northern Territory; 3 in Western Australia and; 3 in South Australia (Figure 1, Table 1 and Table 2). From the commencement of the outbreak to 30 June 2018[‡], there were a total of 2175 outbreak cases (category 1 and 2 cases) reported across the four outbreak affected jurisdictions: 1170 in Queensland (from January 2011); 802 in the Northern Territory (from July 2013); 165 in Western Australia (from June 2014), and; 38 in South Australia (from November 2016) (Table 1 and 2).

Figure 1. Epidemic curve showing category 1 infectious syphilis[‡] outbreak cases notified in Aboriginal and Torres Strait Islander people residing in affected regions[‡] of Queensland, the Northern Territory, Western Australia and South Australia from commencement of the outbreak in each jurisdiction to 30 June 2018^{‡,y}




Table 1. Characteristics of category 1 infectious syphilis^α outbreak cases notified in Aboriginal and Torres Strait Islander people residing in affected regions^β of Queensland, the Northern Territory, Western Australia and South Australia, to 30 June 2018^{‡,‡}

	North Qld (four HHSs ^β)	NT (seven regions ^β)	WA (Kimberley region)	SA (two regions ^β)
Situation to-date, 30 June 2018				
Outbreak commencement month/year	January 2011	July 2013	June 2014	November 2016
Total number of cases ^α	1144	796	165	38
% Male / % Female	47% / 53%	46% / 54%	36% / 64%	45% / 55%
% 15-29 year age group	67%	64%	71%	45%
Congenital cases, confirmed (probable) ^α	4 (4)	2 (2)	0 (0)	1 (0)
-number of deaths in congenital cases, confirmed (probable)	3 (3)	0 (0)	0 (0)	0 (0)
Last reporting month, 1 – 30 June 2018				
Number of cases ^α	22	26	3	3
% Male / % Female	32% / 68%	54% / 46%	33% / 67%	67% / 33%
% 15-29 year age group	59%	27%	67%	33%

Table 2. Number of category 2 infectious syphilis^α cases, to 30 June 2018^{‡,‡}

	North Qld (four HHSs ^β)	NT (seven regions ^β)	WA (Kimberley region)	SA (two regions ^β)
Situation to-date, 30 June 2018				
Outbreak commencement month/year	January 2011	July 2013	June 2014	November 2016
Aboriginal and Torres Strait Islander people [§]	11	2	-	-
Non-Indigenous people ^μ	15	4	-	-
Last reporting month, 1 -30 June 2018				
Aboriginal and Torres Strait Islander people [§]	-	-	-	-
Non-Indigenous people ^μ	-	-	-	-

Notes:

[‡] 'Diagnosis date' was used to define the period of analysis. This date represents either the onset date or where the date of onset was not known, the earliest of the specimen collection date, the notification date, or the notification receive date.

^α Cases defined as per the MJSO syphilis outbreak case definition:

Nationally, an infectious syphilis outbreak case is defined as: any person who is newly diagnosed with confirmed or probable infectious syphilis according to the CDNA national surveillance case definition for infectious syphilis, **AND**, is an Aboriginal or Torres Strait Islander person who resides in any of the following outbreak declared regions as defined and documented by that jurisdiction, at or after the dates indicated: **Qld** - Torres and Cape Hospital and Health Service area (from 1 December 2012); Cairns and Hinterland Hospital and Health Service area (from 1 August 2013); North West Hospital and Health Service area (from 1 January 2011); Townsville Hospital and Health Service area (from 1 January 2014); **NT** - Alice Springs or Barkly district (from 1 July 2013); Katherine district (from 1 May 2014); East Arnhem district (from 1 November 2015); Darwin Rural and Urban (from 1 January 2017); **WA** - Kimberley region, Western Australia (from 1 June 2014), **SA** - Far North and Western and Eyre regions (from 15 November 2016) (category 1 outbreak cases) **OR**, is a sexual contact of a confirmed outbreak case (category 2 outbreak cases).

^β Affected regions include Torres and Cape, Cairns and Hinterland, North West, and Townsville Hospital and Health Services in Queensland; Alice Springs Urban, Alice Springs Rural, Barkly, East Arnhem, Katherine, Darwin Urban and Darwin Rural regions in the Northern Territory; the Kimberley health region in Western Australia, and; Far North and Western and Eyre regions in South Australia.

^α Cases defined as per the MJSO syphilis outbreak case definition: any newly diagnosed case of confirmed or probable congenital syphilis that meets the CDNA national surveillance case definition whose mother's syphilis infection prior to or during pregnancy meets the infectious syphilis outbreak case definition.

[§] Aboriginal and Torres Strait Islander people who are sexual contacts of a confirmed outbreak case and reside outside an outbreak declared region at the time of diagnosis.

^μ Non-Indigenous people who are sexual contacts of a confirmed outbreak case and reside in or out of an outbreak declared region at the time of diagnosis.

[‡] Please note that **all data are provisional** and subject to change due to ongoing case investigation.