

6. Robbins JB, Lee C-J, Rastogi SC *et al.* Comparative immunogenicity of group 6 pneumococcal type 6A (6) and type 6B (26) capsular polysaccharides. *Infect Immun* 1979;26:1116-1122.
7. Nielsen SV, Henrichsen J. Capsular types of *Streptococcus pneumoniae* isolated from blood and CSF during 1982-1987. *Clin Infect Dis* 1992;15:794-798.
8. Butler JC, Breiman R, Lipman HB. *et al.* Serotype distribution of *Streptococcus pneumoniae* infections among preschool children in the United States, 1978-1994: implications for the development of a conjugate vaccine. *J Infect Dis* 1995;171:885-889.
9. Eskola J, Takala A, Kela E. *et al.* Epidemiology of invasive pneumococcal infections in children in Finland. *JAMA* 1992;268:3323-3327.
10. Saha SK, Rikitomi N, Biswas D. *et al.* Serotypes of *Streptococcus pneumoniae* causing invasive childhood infections in Bangladesh, 1992 to 1995. *J Clin Microbiol* 1997;35:785-787.
11. Barker J, Gratten M, Riley I. *et al.* Pneumonia in children in the Eastern Highlands of Papua New Guinea: a bacteriologic study of patients selected by standard clinical criteria. *J Infect Dis* 1989;159:348-352.
12. Kallenius G, Hedlund J, Svenson SB. *et al.* Pneumococcal bacteraemia in Sweden. *Lancet* (letter) 1997; 1:1910.
13. Austrian R. Epidemiology of pneumococcal capsular types causing pediatric infections. *Pediatr Infect Dis J* 1989;8:S21-S22.
14. Siegel J, Poziviak CS, Michaels RH. Serotypically defined pneumococcal infections in children. *J Pediatr* 1978;93:249-250.
15. Scott JAG, Hall AJ, Ragan R. *et al.* Serogroup-specific epidemiology of *Streptococcus pneumoniae*: associations with age, sex, and geography in 7000 episodes of invasive disease. *Clin Infect Dis* 1996;22:973-981.
16. Paton J, Toogood IR, Cockington RA. *et al.* Antibody response to pneumococcal vaccine in children aged 5 to 15 years. *Am J Dis Child* 1986;140:135-138.
17. Heffron R. Pneumonia with special reference to pneumococcus lobar pneumonia. 2nd printing. Cambridge, Massachusetts: Harvard University Press, 1979.
18. Finland M, Barnes MW. Changes in occurrence of capsular serotypes of *Streptococcus pneumoniae* at Boston City Hospital during selected years between 1935 and 1974. *J Clin Microbiol* 1977;5:154-166.
19. Barry MA, Craven DE, Finland M. Serotypes of *Streptococcus pneumoniae* isolated from blood cultures at Boston City Hospital between 1979 and 1982. *J Infect Dis* 1984;149:449-452.
20. Lund E. Types of pneumococci found in blood, spinal fluid and pleural exudate during a period of 15 years (1954-1969). *Acta Pathol Microbiol Scand (B)* 1970;78:333-336.
21. Morch E. On the frequency of pneumococcus types in Denmark 1939-1947. *Acta Pathol Microbiol Scand* 1949;26:83-92.
22. Douglas RM, Paton JC, Duncan SJ. *et al.* Antibody response to pneumococcal vaccination in children younger than five years of age. *J Infect Dis* 1983;148:131-137.
23. Giebink GS. Preventing pneumococcal disease in children: recommendations for using the current pneumococcal vaccine. *Pediatr Infect Dis J* 1985;4:343-348.
24. Sniadack DH, Schwartz B, Lipman H. *et al.* Potential interventions for the prevention of childhood pneumonia: geographic and temporal differences in serotype and serogroup distribution of sterile site pneumococcal isolates from children. *Pediatr Infect Dis J* 1995;14:5-10.

Australian Recommendations for the Influenza Vaccine Composition for the 1999 Season

The meeting of the Australian Influenza Vaccine Committee (AIVC) on Influenza Vaccines was convened on 7 October 1998, in order to select virus strains for the manufacture of Influenza Vaccine for 1999 Season.

Having considered the information on international surveillance by WHO and up-to-date epidemiology and strains characterisation presented at the meeting the Committee considered that the WHO recommendations on the composition of vaccines for 1999 Season should be followed:

- **A (H3N2)** an A/Sydney/5/97 (H3N2)-like strain, 15 µg HA per dose.
- **A (H1N1)** an A/Beijing/262/95 (H1N1)-like strain, 15 µg HA per dose.
- **B** a B/Beijing/184/93- like strain, 15 µg HA per dose.

It was also determined that the following viruses are suitable vaccine strains:

- The high yield reassortant viruses IVR-108 and RESVIR-13 are A/Sydney/5/97 (H3N2)-like strains
- The high yield reassortant virus X-127 is accepted as an A/Beijing/262/95 (H1N1)-like strain
- B/Harbin/7/94 is accepted as a B/Beijing/184/93-like strain