

Australian Respiratory Surveillance Report

Key messages

This report presents a national epidemiological update for acute respiratory infections, including coronavirus disease 2019 (COVID-19), influenza and respiratory syncytial virus (RSV), with a focus on the current reporting period (11 August to 24 August 2025) and earlier severity reporting periods (up to 10 August 2025).

In the community: In the last fortnight, influenza-like illness among national helpline callers and the rate of helpline callers referred to seek urgent medical care both increased compared to the previous fortnight. Self-reported new fever and cough symptoms among community survey participants also increased slightly compared to the previous fortnight; however, slightly fewer community survey participants reported taking time off work due to respiratory illness. COVID-19 cases continued to decrease in the last fortnight and remain lower than at the same time in previous years. Influenza cases decreased last fortnight; however, national influenza case numbers were higher than observed at the same time in previous years. This may indicate that case numbers are decreasing more slowly than observed in previous seasons, which may potentially be driven by higher case numbers in school aged children (5–19 years) currently, but it is too early to definitely determine this trend. RSV cases continued to decrease in the last fortnight, continuing the trend in national case numbers observed since mid-July 2025.

In general practice: In the last fortnight, there were fewer general practice consultations with influenza-like illness (defined as new fever and cough symptoms) at sentinel surveillance sites than in the previous fortnight. Since late June 2025, influenza-like illness consultation rates have exceeded the five-year average but remain slightly lower than observed at the same time last year.

In hospitals: Sentinel hospital-based surveillance indicates that admissions with severe acute respiratory infections have been decreasing since a peak in late June 2025; however, admissions with influenza among school aged children (5–16 years) have continued to increase recently. The proportion of patients who were admitted directly to intensive care at a sentinel hospital site remains low. At sentinel hospitals, more children (those aged 16 years and younger) were admitted with RSV than with influenza or COVID-19. In contrast, more adults were admitted with COVID-19 compared to influenza or RSV. Sentinel intensive care surveillance shows that intensive care admissions with severe acute respiratory infections have been declining since late June 2025. In June and July 2025, most patients were admitted with influenza. A higher proportion of intensive care admissions with influenza and parainfluenza required invasive mechanical ventilation. The duration of intensive care stay is relatively similar between illnesses. In the last fortnight, both the average number of COVID-19 cases occupying intensive care beds and the average number of intensive care staff unavailable due to illness decreased.

Deaths: COVID-19 has remained the leading cause of acute respiratory infection mortality across 2023–2025. All three of these acute respiratory infections are more likely to cause death in older age groups than younger age groups.

In laboratories: Test positivity for SARS-CoV-2 and influenza decreased in the last fortnight, while test positivity for RSV increased slightly. The SARS-CoV-2 variant under monitoring, NB.1.8.1, is the dominant SARS-CoV-2 variant in the last 28 days (14 July to 10 August 2025) accounting for 72.2% of sequences in Australia. Small numbers of sequences of other variants under monitoring, including XFG, LP.8.1, KP.3.1.1 and XEC continue to be observed in Australia.

Vaccine coverage, effectiveness and match: Nationally, fewer adults have received a COVID-19 vaccine in the last 12 months compared to the 12 months prior. Influenza vaccine coverage is 30.2% this year, lower than at the same time in 2023 or 2022. Since the commencement of the National RSV Mother and Infant Protection Program, 109,939 Abrysvo doses have been administered. In the last six months, nirsevimab uptake is 21.0% nationally. Of influenza isolates characterised in 2025 thus far, over 98% have been a good match to the corresponding 2025 vaccine components.

Australian Respiratory Surveillance Report

This report was prepared by Lauren Welsh, Suzie Whitehead, Algreg Gomez and Jenna Hassall on behalf of the interim Australian Centre for Disease Control. We thank the staff and participants from the surveillance systems who contribute data for acute respiratory illness surveillance across Australia.

The report presents a national overview of acute respiratory infections in Australia, drawing information from several different surveillance systems. These surveillance systems help us to understand the distribution of acute respiratory illnesses in the community, the severity of infections including which populations might be at risk, and the impact of acute respiratory illnesses on the community and health system in Australia.

Surveillance indicators presented in this report are based on the [Australian National Surveillance Plan for COVID-19, Influenza, and RSV](#). Please refer to the [Technical Supplement – Australian Respiratory Surveillance Report](#) for information on our surveillance sources and data considerations, including the considerable impact of the COVID-19 pandemic on acute respiratory infection surveillance in Australia. A summary of data considerations for this report are provided below:

- Due to the dynamic nature of the surveillance systems used in this report, surveillance data are considered preliminary and subject to change as updates are received, with the most recent weeks considered particularly incomplete. Data in this report may vary from data reported in other national reports and reports by states and territories.
- Data in this report are presented by date of event (diagnosis, admission or death) or by the International Organization for Standardization (ISO) week date system, with weeks defined as seven-day periods which begin on a Monday and end on a Sunday. The ISO week date system is used to support trends comparisons over time more effectively. The current reporting period includes 11 August to 24 August 2025 and where comparisons to the previous fortnight are made, this includes 28 July to 10 August 2025.
- In Australia, states and territories (the Australian Capital Territory [ACT], New South Wales [NSW], the Northern Territory [NT], Queensland [Qld], South Australia [SA], Tasmania [Tas], Victoria [Vic] and Western Australia [WA]) report notified cases to the **National Notifiable Diseases Surveillance System (NNDSS)** based on the [Australian national surveillance case definitions](#). NNDSS data are analysed and reported based on diagnosis date, which is the true onset date of a case if known, otherwise it is the earliest of the specimen date, the notification date or the notification received date. The NNDSS data for this report were extracted on 28 August 2025.
- To account for the lag in collection and provision of severity data from some surveillance systems, and for the time delay between illness onset and the development of severe disease outcomes, cases with an admission date or a diagnosis date in the last two weeks are excluded from severity analyses for hospitalisations and intensive care admissions. As such, the severity reporting periods are two weeks behind the end of the current reporting period. For this report, severity reporting includes data from 28 July to 10 August 2025 unless specified otherwise. Where comparisons to the previous severity fortnight are made this includes 14 July to 27 July 2025.
- Death registrations from the Australian Bureau of Statistics (ABS) Provisional Mortality Statistics are now used as the primary data source for measuring acute respiratory infection associated deaths. The ABS mortality data is sourced from the Registry of Births, Deaths and Marriages and is separate from the NNDSS. Registration-based mortality data needs time to be received and processed. For this reason, mortality statistics in this report may lag by at least two months.
- Analysis and reporting outputs were produced using R Statistical Software v4.3.1. While every care has been taken in preparing this report, the Australian Government Department of Health, Disability and Ageing does not accept liability for any injury or loss or damage arising from the use of, or reliance upon, the content of the report or Technical Supplement. For further information about this report refer to the [Technical Supplement – Australian Respiratory Surveillance Report](#) or contact respiratory-surveillance@health.gov.au.


Community surveillance

Community surveillance monitors respiratory illnesses in the community, providing information on the number of people reporting respiratory symptoms, testing practices, and the impact of respiratory illnesses.

Community surveillance includes notification data obtained from laboratory tests for infections. Infections that are diagnosed and notified are only a subset of the total number of infections occurring in the community.

- In the last fortnight (11 August to 24 August 2025), there were slightly more Healthdirect helpline callers with influenza-like illness (407 per 1,000 callers per fortnight) than in the previous fortnight (400 per 1,000 callers per fortnight) (Figure 1).
- Since late June 2025 the rate of influenza-like illness among helpline callers has been higher than the rate of influenza-like illness at the same time in 2024 and 2023 (Figure 1).

Figure 1: Rate of influenza-like illness per 1,000 helpline callers by year and week of call*, Australia†, 22 March 2023 to 24 August 2025




Source: Healthdirect Australia

* Healthdirect data prior to 22 March 2023 are unavailable as prior to this date a different data collection method was used.

† The Healthdirect helpline operates in all states and territories except Qld; therefore influenza-like illness trends will not be representative of Qld and may be underrepresented. See the [Technical Supplement](#) for more information.

- In the last fortnight, there were more Healthdirect helpline callers with influenza-like illness referred to seek urgent medical care (210 per 1,000 callers per fortnight) than in the previous fortnight (198 per 1,000 callers per fortnight) (Figure 2).
 - Callers referred to seek urgent medical care include those referred to call triple zero, attend a hospital emergency department, contact a virtual emergency department, urgent care clinic or see a general practitioner within two hours.
- In the last fortnight, referral pathways for influenza-like illness varied across Australian jurisdictions. NSW, SA and Vic had the highest proportion of callers referred to see a general practitioner (GP) or seek other urgent medical care (Figure 2). By comparison, the ACT, the NT and Tas had a higher proportion of callers who were recommended to attend a hospital emergency department (Figure 2). These differences may reflect variations in healthcare access and service models, and should be interpreted with caution.

Figure 2: Proportion of referral categories* for helpline callers with influenza-like illness by jurisdiction† and call date, Australia, 1 January to 24 August 2025




Source: Healthdirect Australia

* See other health professionals category includes pharmacist, dentist, mental health provider, primary maternity care, poison information centre or other.

† The Healthdirect helpline operates in all states and territories except Qld; therefore influenza-like illness referral trends are not provided for Qld. See the [Technical Supplement](#) for more information.

- In the last fortnight, slightly more FluTracking survey participants reported new fever and cough symptoms (2.0%), than in the previous fortnight (1.9%) (Figure 3).
- In the last fortnight, more survey participants with new fever and cough symptoms used a rapid antigen test (RAT) (52.7%; 601/1,141) than a polymerase chain reaction (PCR) test (11.0%; 125/1,141) to test for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).
- Self-reported SARS-CoV-2 RAT positivity was slightly lower in the last fortnight (14.8%; 89/601) than in the previous fortnight (15.5%; 108/695). Self-reported SARS-CoV-2 PCR positivity was lower in the last fortnight (8.0%; 10/125) than in the previous fortnight (12.6%; 17/135).
- In the last fortnight, 10.3% (117/1,141) of survey participants with new fever and cough symptoms used a PCR test to test for influenza. Self-reported influenza PCR positivity was lower this fortnight (32.5%; 38/117), than in the previous fortnight (36.0%; 45/125).
- In the last fortnight, fewer survey participants reported taking three or more days off work or normal duties due to fever and cough symptoms (45.4%; 518/1,141), than in the previous fortnight (51.3%; 631/1,229).
- In the last fortnight, for the first time in 2025, the weekly percentage of FluTracking participants reporting new fever and cough symptoms was higher than observed at the same time in previous years and the five-year average (Figure 3). However, the weekly percentage remained relatively similar to the trends observed at the same time in 2022, 2024 and the five-year average (Figure 3).

Figure 3: Age standardised percentage of survey participants reporting new fever and cough symptoms compared with the five-year average* by year and week of survey, Australia, 2022 to 24 August 2025



Source: FluTracking

* From 2020, FluTracking expanded their data capture period to year-round. Data before May and after October for any year before 2020 are not available for historical comparisons. The years 2020 and 2021 are excluded when comparing the current season to historical periods when influenza virus has circulated without public health restrictions. As such, the five-year average includes the years 2018 to 2019 and 2022 to 2024. Please refer to the [Technical Supplement](#) for interpretation of the five-year average.

- In the last fortnight (11 August to 24 August 2025), there was an 8.2% decrease in COVID-19 cases, a 6.5% decrease in influenza cases, and a 6.8% decrease in RSV cases.

Table 1: Notified cases and notification rate per 100,000 population by disease, five-year age group, and jurisdiction*†, Australia, 1 January to 24 August 2025

	COVID-19			Influenza			RSV		
	Reporting period (n)	Year to date (n)	Year to date (rate)	Reporting period (n)	Year to date (n)	Year to date (rate)	Reporting period (n)	Year to date (n)	Year to date (rate)
Age group (years)									
0–4	840	13,933	923	4,076	37,231	2,468	4,157	65,682	4,353
5–9	374	3,945	245	5,760	44,646	2,771	1,016	10,138	629
10–14	386	4,313	258	4,573	30,092	1,797	535	4,265	255
15–19	330	4,742	285	2,480	18,042	1,085	304	2,854	172
20–24	277	4,999	279	1,256	10,404	581	211	2,312	129
25–29	338	6,156	308	1,143	10,107	506	260	2,841	142
30–34	428	7,538	370	1,724	14,873	729	283	3,613	177
35–39	497	8,628	435	2,366	20,071	1,011	293	3,681	185
40–44	485	8,368	452	2,335	20,461	1,105	294	3,177	172
45–49	389	7,131	438	1,786	15,848	973	268	2,912	179
50–54	395	7,180	425	1,548	14,411	853	337	3,572	211
55–59	340	6,981	455	1,504	13,244	864	345	3,864	252
60–64	399	7,447	485	1,363	12,802	834	412	4,412	288
65–69	376	7,640	562	1,288	11,319	833	417	4,649	342
70+	2,069	47,745	1,429	4,036	37,133	1,112	2,039	19,289	577
Jurisdiction									
ACT	112	2,317	489	753	6,334	1,336	246	2,466	520
NSW	2,989	66,240	781	14,155	119,663	1,410	3,086	60,098	708
NT	76	1,063	417	227	3,221	1,263	40	677	265
Qld	2,047	30,697	550	8,850	62,526	1,119	1,830	24,768	443
SA	430	8,930	475	2,535	22,188	1,181	1,663	8,541	455
Tas	153	2,158	375	673	4,643	807	253	1,769	307
Vic	1,606	25,173	361	7,041	71,067	1,018	2,539	30,277	434
WA	527	10,313	348	3,022	21,260	717	1,516	8,678	293
Total	7,940	146,891	540	37,256	310,902	1,143	11,173	137,274	505


Source: National Notifiable Diseases Surveillance System (NNDSS). Notification data for RSV are incomplete for Tas from 22 August 2025.

* Rate per 100,000 population for the given time period. Population data are based on the Australian Bureau of Statistics (ABS) [Estimated Resident Population \(ERP\) for the reference period June 2024, released 12 December 2024](#).

† Total includes cases with missing age.


- In the last fortnight, the number of COVID-19 cases reported each week remained relatively stable; however, overall case numbers have continued to decrease since reaching a recent peak in late June 2025 (Figure 4).
- The number of COVID-19 cases this year to date (n=146,891) is 31.6% less than the number of cases observed in the same time period last year (n=214,782) (Figure 4).
- In the last fortnight, COVID-19 notification rates remained relatively stable across most jurisdictions; however, notification rates decreased in NSW, Qld and SA, and increased slightly in the NT compared with the previous fortnight (Figure 5).
- In the year to date, COVID-19 notification rates remain highest in people aged 70 years or over, likely due to higher case ascertainment from targeted testing strategies for populations at-risk of severe disease or who live in a high-risk setting such as a residential aged care home (Table 1).
- In the year to date, COVID-19 notification rates are highest in NSW and lowest in WA (Table 1).

Figure 4: Notified COVID-19 cases (laboratory-confirmed only) by year and week of diagnosis, Australia, 2022 to 24 August 2025



Source: National Notifiable Diseases Surveillance System (NNDSS)


Figure 5: Notification rates* per 100,000 population for COVID-19 cases by state or territory and week of diagnosis, Australia, 1 January to 24 August 2025



Source: National Notifiable Diseases Surveillance System (NNDSS)

* Rate per 100,000 population for the given time period. Population data are based on the Australian Bureau of Statistics (ABS) [Estimated Resident Population \(ERP\)](#) for the reference period June 2024, released 12 December 2024

Figure 6: Notified influenza cases and five-year average* by year and week of diagnosis, Australia, 2022 to 24 August 2025




Source: National Notifiable Diseases Surveillance System (NNDSS)

* The years 2020 and 2021 are excluded when comparing the current season to historical periods when influenza virus has circulated without public health restrictions. As such, the five-year average includes the years 2018 to 2019 and 2022 to 2024. Please refer to the [Technical Supplement](#) for interpretation of the five-year average.

- In the last fortnight, the number of influenza cases nationally decreased; however, influenza case numbers in the last fortnight were higher than observed at the same time in previous years (Figure 6). This may indicate that case numbers are decreasing more slowly than observed in previous seasons, which may potentially be driven by higher case numbers in school aged children currently, but it is too early to definitely determine this trend.
- The number of influenza cases this year to date (n=310,902) is 1.9% less than the number of cases observed in the same time period last year (n=316,894); however, case numbers currently remain elevated so this trend may change in future (Figure 6).
- In the last fortnight, influenza notification rates decreased across most jurisdictions except for SA, Tas and WA where notification rates increased compared with the previous fortnight (Figure 7).

- In the last fortnight, the number of influenza cases among school aged children (5–19 years) remained elevated, following an increase in cases among this age group in the previous fortnight (11 August to 24 August 2025). This trend is likely due to increased population mixing following the school holiday period (Figure 8). The number of influenza cases among other age groups decreased during this time (Figure 8).
- In the year to date, influenza notification rates remain highest in children aged 5–9 years and children aged 0–4 years (Table 1).
- In the year to date, influenza notification rates are highest NSW and lowest in WA (Table 1).

Figure 7: Notification rates* per 100,000 population for influenza cases by state or territory and week of diagnosis, Australia, 1 January to 24 August 2025




Source: National Notifiable Diseases Surveillance System (NNDSS)

* Rate per 100,000 population for the given time period. Population data are based on the Australian Bureau of Statistics (ABS) Estimated Resident Population (ERP) for the reference period June 2024, released 12 December 2024.

- In the last fortnight, most influenza notifications were influenza A(Unsubtyped) (60.5%; 22,540/37,256), followed by influenza B (37.1%; 13,804/37,256), then influenza A(H1N1) (1.8%; 665/37,256), and influenza untyped (0.4%; 142/37,256). In the last fortnight, there were 48 influenza A&B co-detections (Figure 8).
- In the year to date, influenza A(Unsubtyped) has accounted for most cases across all age groups, followed by influenza B. The proportion of influenza B cases is the highest in the 5–19 years age group. There has been a small number of influenza A(H1N1) and influenza A(H3N2) cases across all age groups (Figure 8).
 - There is a comparatively higher proportion of influenza B cases this season than observed in 2024. While influenza B is often a good match with the seasonal influenza vaccine strain, influenza B can result in more severe infections in children.
- In the year to date, influenza A(Unsubtyped) has accounted for the majority of influenza cases across all jurisdictions. Several jurisdictions have been experiencing increasing numbers of influenza B cases in the year to date; however, the proportion of influenza B and influenza A varies week-on-week (Figure 9).
- Influenza A(H1N1) and influenza A(H3N2) cases were most commonly observed in Tas and WA (Figure 9); however, trends in influenza subtypes should be interpreted with care as there are jurisdictional differences in the number and selection of influenza samples that undergo typing.


Figure 8: Notified influenza cases by influenza subtype, age group*, and week of diagnosis, Australia, 1 January to 24 August 2025



Source: National Notifiable Diseases Surveillance System (NNDSS)

* Axis varies between age groups.

Figure 9: Notified influenza cases by influenza subtype, jurisdiction*, and week of diagnosis, Australia, 1 January to 24 August 2025




Source: National Notifiable Diseases Surveillance System (NNDSS)

* Axis varies between jurisdictions.

- In the last fortnight, the number of RSV cases decreased, continuing the gradually decreasing trend in national case numbers observed since mid-July 2025 (Figure 10).
- The number of RSV cases this year to date (n=137,274) is 6.4% less than the number of cases observed in the same time period last year (n=146,681), largely due to comparatively fewer cases across May and June 2025 (Figure 10).
- In the last fortnight, RSV notification rates decreased or remained relatively stable across all jurisdictions (Figure 11).
- In the year to date, RSV notification rates remain considerably higher in children aged 0–4 years than in other age groups (Table 1).
- In the year to date, RSV notification rates are highest in NSW and lowest in the NT (Table 1).


Figure 10: Notified RSV cases by year and week of diagnosis*, Australia, 2023 to 24 August 2025



Source: National Notifiable Diseases Surveillance System (NNDSS)

* RSV became notifiable in all states and territories on 1 September 2022 and comprehensive national notification data became available after this point. For this reason, RSV notification trends are only presented from 1 January 2023.

Figure 11: Notification rates* per 100,000 population for RSV cases by state or territory and week of diagnosis, Australia, 1 January to 24 August 2025



Source: National Notifiable Diseases Surveillance System (NNDSS). Notification data for RSV are incomplete for Tas from 22 August 2025.


* Rate per 100,000 population for the given time period. Population data are based on the Australian Bureau of Statistics (ABS) [Estimated Resident Population \(ERP\)](#) for the reference period June 2024, released 12 December 2024.

Primary care surveillance

Primary care surveillance monitors the number and characteristics of people who have presented to a general practice with influenza-like illness and provides insight on the different respiratory pathogens that are causing illness in the community.

- In the last fortnight (11 August to 24 August 2025), there were fewer general practice consultations for influenza-like illness (9.9 notifications per 1,000 consultations per fortnight) compared to fortnight (11.1 notifications per 1,000 consultations per fortnight) (Figure 12).
- From late June 2025, influenza-like illness consultation rates have been higher than in 2022, 2023 and the five-year average but remain slightly lower than observed at the same time in 2024 (Figure 12).

Figure 12: Rate of influenza-like illness notifications per 1,000 consultations per week in sentinel general practice sites compared with the five-year average by year and week of consultation*†, Australia, 2022 to 24 August 2025




Source: Australian Sentinel Practice Research Network (ASPREN)

* The years 2020 and 2021 are excluded when comparing the current season to historical periods when influenza virus has circulated without public health restrictions. As such, the five-year average includes the years 2018 to 2019 and 2022 to 2024. Please refer to the [Technical Supplement](#) for interpretation of the five-year average.

† Please refer to the [Technical Supplement](#) for notes on impact of COVID-19 on ASPREN data.

- In the last fortnight, 58.2% (82/141) of people attending general practice with influenza-like illness who were tested have then tested positive for a respiratory pathogen.
- In the last fortnight, influenza (34.1%; 28/82) was the most commonly detected pathogen, followed by rhinovirus (23.2%; 19/82) and RSV (15.9%; 13/82) (Figure 13).
- In the year to date, 66.9% (1,331/1,989) of people attending general practice with influenza-like illness who were tested have then tested positive for a respiratory pathogen.
- In the year to date, rhinovirus (33.9%; 451/1,331) has been the most commonly detected pathogen, followed by influenza (30.0%; 399/1,331), RSV (9.7%; 129/1,331), SARS-CoV-2 (9.7%; 129/1,331) and adenovirus (4.7%; 63/1,331) (Figure 13).

Figure 13: Number of samples tested for respiratory pathogens among people with influenza-like illness attending sentinel general practice sites by respiratory pathogen and week of specimen collection, Australia, 1 January to 24 August 2025



Source: Australian Sentinel Practice Research Network (ASPREN)


Note: All ASPREN swab samples are transported to the SA Pathology laboratory in Adelaide to be tested for viral and bacterial respiratory pathogens via a multiplex real-time reverse transcription polymerase chain reaction (RT-PCR) assay using in-house primers.

Hospital-based surveillance

Hospital-based surveillance monitors persons with more severe illness who have been admitted to hospital for their respiratory illness (severe acute respiratory infections). Hospital-based surveillance also measures the ability of the health system to cope with the number of severe acute respiratory infection admissions to ensure delivery of safe, timely and quality health care.

- In the last severity reporting period (28 July to 10 August 2025), fewer patients were admitted to a sentinel hospital with a severe acute respiratory infection (n=738), than in the previous severity reporting period (n=922).
 - In the last severity reporting period, at sentinel hospitals there was 24.6% fewer admissions with COVID-19 (from 118 to 89), 17.0% fewer admissions with influenza (from 495 to 411), and 23.0% fewer admissions with RSV (from 309 to 238), compared to the previous severity reporting period.
 - In the last severity reporting fortnight, at sentinel hospitals admissions with influenza among school aged children continued to increase despite an overall decrease in the number of admissions with influenza (Figure 15)
- In the year to date for severity reporting (1 January to 10 August 2025), there have been 9,641 admissions with severe acute respiratory infections at sentinel hospitals. Most patients with a severe acute respiratory infection have been admitted with influenza (n=4,120) followed by RSV (n=3,133) (Figure 14).

Figure 14: Total number of patients (children and adults) admitted with a severe acute respiratory infection to sentinel hospitals by disease and week of admission, Australia, 1 January to 10 August 2025




Source: Influenza Complications Alert Network (FluCAN)

- Patients admitted to sentinel hospitals with influenza have mostly been admitted with influenza A (80.0%; 3,295/4,120), while 19.6% (807/4,120) were admitted with influenza B.
 - Most hospital admissions with influenza A have been with influenza A(Unsubtyped) (68.0%; 2,239/3,295), followed by influenza A(H1N1) (30.7%; 1,012/3,295) and then influenza A(H3N2) (1.3%; 44/3,295).
- In the year to date for severity reporting, influenza A was the most commonly detected influenza type in all age groups. Influenza A(H1N1) and influenza A(H3N2) subtypes have been more commonly observed in adults than children, while influenza B has been more commonly observed in children. Of

note, school aged children (5–16 years) had the highest proportion of influenza B compared with influenza A (Figure 15).

- While influenza B is often a good match with the seasonal influenza vaccine strain, influenza B can result in more severe infections in children.
- Trends in influenza subtypes should be interpreted with care as there may be differences in the number and selection of influenza samples that undergo typing.

Figure 15: Number of patients admitted with influenza to sentinel hospitals by influenza subtype, age group*, and week of admission, Australia, 1 January to 10 August 2025



Source: Influenza Complications Alert Network (FluCAN)

* Axis varies between age groups. The age distribution of admissions with influenza may not reflect the age distribution of all patients.

- In the year to date for severity reporting, more children (those aged 16 years and younger) have been admitted to sentinel hospitals with RSV than with influenza or COVID-19 (Table 2a).
- Children admitted to sentinel hospitals with influenza tended to be older than children admitted with COVID-19 or RSV (Table 2a).
- Children admitted to sentinel hospitals with RSV had a slightly longer length of hospital stay compared to children with influenza or COVID-19; however, the difference in the length of stay was minor. The proportion of children admitted directly to intensive care was slightly higher for COVID-19 than influenza and RSV (Table 2a).

- Sadly, there have been a small number of children admitted with a severe acute respiratory infections who have died in sentinel hospitals (Table 2a).

Table 2a: Demographic characteristics and outcomes for children admitted with a severe acute respiratory infection to a sentinel hospital by disease*†‡, Australia, 1 January to 10 August 2025

	COVID-19	Influenza	RSV
	Year to date for severity reporting (n=750)	Year to date for severity reporting (n=2,287)	Year to date for severity reporting (n=2,664)
Age (years)			
Median [IQR]	1 [0–4]	4 [1–8]	1 [0–2]
Age group (years)			
< 6 months	241 (32.1%)	140 (6.1%)	633 (23.8%)
6 months – 4 years	335 (44.7%)	1073 (46.9%)	1838 (69.0%)
5–16 years	174 (23.2%)	1074 (47.0%)	193 (7.2%)
Indigenous status			
Aboriginal and Torres Strait Islander	71 (9.5%)	152 (6.6%)	160 (6.0%)
Length of hospital stay (days)†			
Median [IQR]	1 [1–3]	1 [1–2]	2 [1–3]
Patient admission location‡			
Admitted to hospital ward	694 (92.5%)	2165 (94.7%)	2523 (94.7%)
Admitted to intensive care directly	56 (7.5%)	122 (5.3%)	141 (5.3%)
Discharge status†			
Alive	549 (73.2%)	1726 (75.5%)	1830 (68.7%)
Died	1 (0.1%)	5 (0.2%)	1 (< 0.1%)
Incomplete/missing	200 (26.7%)	556 (24.3%)	833 (31.3%)

Source: Influenza Complications Alert Network (FluCAN)

* Does not include patients with missing age; therefore, the sum of age-specific totals above may not equal the total number of patients.

† For patients who are still in hospital data may not be complete; therefore, these data are not included in the length of stay or discharge status. In addition, length of stay data excludes patients that acquired their infection in hospital.

‡ Admission location reflects the initial admission ward. Some patients may be initially admitted to general ward then later admitted to an intensive care and this is not reflected here. Does not include patients with missing admission location; therefore, the sum of admission location specific totals above may not equal the total number of patients.

The Paediatric Active Enhanced Disease Surveillance (PAEDS) network carries out enhanced sentinel hospital surveillance for some acute respiratory infections or conditions in children. PAEDS data for acute respiratory infections in children are presented in the Australian Respiratory Surveillance Reports in the sentinel hospital data from FluCAN. For additional information on [COVID-19 in children](#), [Paediatric Inflammatory Multisystem Syndrome \(PIMS-TS\) following COVID-19](#), [influenza in children](#), or [RSV in children](#) please visit the [PAEDS](#) webpages and dashboards.

- In the year to date for severity reporting, more adults (those aged 17 years and over) have been admitted to sentinel hospitals with influenza than with COVID-19 or RSV (Table 2b).
- Adults admitted to sentinel hospitals with COVID-19 or RSV were predominately 65 years and over, whereas the proportion of admissions with influenza was only slightly higher in the 65 years and over age group compared to the 17–64 years age group (Table 2b).
- Adults admitted to sentinel hospitals with COVID-19 had a slightly longer length of hospital stay compared to adults with influenza or RSV; however, the difference in the length of stay was minor. A higher proportion of adults with influenza were admitted directly to intensive care, compared to adults admitted with COVID-19 or RSV (Table 2b).
- Sadly, there have been a number of adults admitted with a severe acute respiratory infections who have died in sentinel hospitals (Table 2b).

Table 2b: Demographic characteristics and outcomes for adults admitted with a severe acute respiratory infection to a sentinel hospital by disease^{†‡}, Australia, 1 January to 10 August 2025**

	COVID-19	Influenza	RSV
	Year to date for severity reporting (n=1,638)	Year to date for severity reporting (n=1,832)	Year to date for severity reporting (n=469)
Age (years)			
Median [IQR]	75 [62–84]	67 [52–77]	73 [61–82]
Age group (years)			
17–64 years	460 (28.1%)	845 (46.1%)	135 (28.8%)
65 years and over	1178 (71.9%)	987 (53.9%)	334 (71.2%)
Indigenous status			
Aboriginal and Torres Strait Islander	92 (5.6%)	113 (6.2%)	29 (6.2%)
Length of hospital stay (days)[†]			
Median [IQR]	5 [2–9]	4 [2–7]	4 [2–8]
Patient admission location[‡]			
Admitted to hospital ward	1547 (94.4%)	1679 (91.6%)	438 (93.4%)
Admitted to intensive care directly	91 (5.6%)	153 (8.4%)	31 (6.6%)
Discharge status[†]			
Alive	1198 (73.1%)	1283 (70.0%)	293 (62.5%)
Died	60 (3.7%)	54 (2.9%)	17 (3.6%)
Incomplete/missing	380 (23.2%)	495 (27.0%)	159 (33.9%)

Source: Influenza Complications Alert Network (FluCAN)


* Does not include patients with missing age; therefore, the sum of age-specific totals above may not equal the total number of patients.

† For patients who are still in hospital data may not be complete; therefore, these data are not included in the length of stay or discharge status. In addition, length of stay data excludes patients that acquired their infection in hospital.

‡ Admission location reflects the initial admission ward. Some patients may be initially admitted to general ward then later admitted to an intensive care and this is not reflected here. Does not include patients with missing admission location; therefore, the sum of admission location specific totals above may not equal the total number of patients.

- Please note, sentinel intensive care data are updated each month, as such the sentinel intensive care surveillance data presented here have not been updated since the previous report.
- In the last severity reporting period for sentinel intensive care (30 June to 27 July 2025), fewer patients have been admitted to a sentinel intensive care with a severe acute respiratory infection (n=432), than in the previous severity reporting period (n=470) (Figure 16). Admissions to sentinel intensive care in the last severity reporting period were predominantly patients admitted with influenza (46.6%; 210/451) (Figure 16).
- In the year to date for severity reporting (1 January to 27 July 2025), most patients were admitted to sentinel intensive care with rhinovirus / enterovirus, followed by influenza (Figure 16; Table 3).


Figure 16: Number of patients admitted with severe acute respiratory infections to a sentinel intensive care by disease and week of admission, Australia, 1 January to 27 July 2025



Source: Short Period Incidence Study of Severe Acute Respiratory Infection (SPRINT-SARI) Australia

Note: A range of diagnostic testing procedures are utilised across hospitals in Australia. SPRINT-SARI does not specify which diagnostic testing method should be utilised as this is the domain of the hospital and treating clinicians. Therefore, virological data from SPRINT-SARI should be interpreted with care.

Figure 17: Number of patients admitted with severe acute respiratory infections to a sentinel intensive care by disease and age group*, Australia, 1 January to 27 July 2025



Source: Short Period Incidence Study of Severe Acute Respiratory Infection (SPRINT-SARI) Australia

Note: 4.9% (113/2,308) of patients had co-infections of respiratory pathogens; therefore, the sum of pathogen-specific totals above may not equal the total number of severe acute respiratory infection patients.

* The age distribution of severe acute respiratory infection intensive care admissions may not reflect the age distribution of all patients.

- In the year to date for severity reporting, admissions to a sentinel intensive care with COVID-19 or influenza have generally been among older people. In contrast, admissions with rhinovirus or RSV have been among younger people, primarily those aged 0–9 years old (Figure 17; Table 3).
- A higher proportion of admissions with influenza and parainfluenza required invasive mechanical ventilation and the length of ventilation was longest among those with influenza. The length of intensive care stay was relatively similar across diseases (Table 3).
- Most patients admitted to a sentinel intensive care with a severe acute respiratory infection have been discharged home. Sadly, a number of patients have died in hospital, predominately among those with COVID-19 (Table 3).

Table 3: Demographic characteristics and outcomes of patients admitted with a severe acute respiratory infection to a sentinel intensive care by disease*†, Australia, 1 January to 27 July 2025

	COVID-19	hMPV	Influenza	Parainfluenza	Rhinovirus	RSV	Other
	Year to date for severity reporting (n=430)	Year to date for severity reporting (n=56)	Year to date for severity reporting (n=687)	Year to date for severity reporting (n=122)	Year to date for severity reporting (n=727)	Year to date for severity reporting (n=297)	Year to date for severity reporting (n=111)
Age (years)							
Median [IQR]	65 [43–75]	35 [4–67]	59 [39–68]	20 [5–67]	12 [4–46]	10 [2–59]	21 [7–59]
Indigenous status							
Aboriginal and Torres Strait Islander	40 (9.3%)	3 (5.4%)	63 (9.2%)	6 (4.9%)	71 (9.8%)	27 (9.1%)	12 (10.8%)
Non-Indigenous	390 (90.7%)	53 (94.6%)	624 (90.8%)	116 (95.1%)	656 (90.2%)	270 (90.9%)	99 (89.2%)
Received invasive mechanical ventilation							
Number (%)	119 (27.7%)	15 (26.8%)	223 (32.5%)	50 (41.0%)	204 (28.1%)	56 (18.9%)	43 (38.7%)
Length of invasive mechanical ventilation (days)*							
Median [IQR]	3 [1–6]	3 [2–9]	5 [2–10]	3 [1–11]	3 [1–7]	3 [1–4]	3 [1–6]
Length of intensive care stay (days)*							
Median [IQR]	3 [2–5]	3 [1–6]	3 [2–7]	2 [1–7]	2 [1–5]	2 [2–4]	3 [2–7]
Length of hospital stay (days)*							
Median [IQR]	7 [4–14]	9 [5–18]	8 [4–15]	7 [3–14]	6 [3–12]	5 [3–9]	8 [4–17]
Patient outcome†							
Ongoing care in intensive care	15 (3.5%)	1 (1.8%)	41 (6.0%)	–	6 (0.8%)	6 (2.0%)	2 (1.8%)
Ongoing care in hospital ward	16 (3.7%)	–	37 (5.4%)	7 (5.7%)	24 (3.3%)	10 (3.4%)	4 (3.6%)
Transfer to other hospital / facility	64 (14.9%)	8 (14.3%)	103 (15.0%)	19 (15.6%)	76 (10.5%)	36 (12.1%)	11 (9.9%)
Discharged home	263 (61.2%)	43 (76.8%)	432 (62.9%)	89 (73.0%)	568 (78.1%)	230 (77.4%)	79 (71.2%)
Died in hospital	70 (16.3%)	4 (7.1%)	72 (10.5%)	6 (4.9%)	48 (6.6%)	15 (5.1%)	14 (12.6%)

Source: Short Period Incidence Study of Severe Acute Respiratory Infection (SPRINT-SARI) Australia


Note: 4.9% (113/2,308) of patients had co-infections of respiratory pathogens; therefore, the sum of pathogen-specific totals above may not equal the total number of severe acute respiratory infection patients.

* For patients receiving ongoing care in intensive care data may not be complete; therefore, data are not included in the length of ventilation or stay.

† Patients who have been admitted with no discharge information for less than 90 days have been assumed to have ongoing care in the hospital. Patients who have no outcome entered or have been admitted for more than 90 days with no discharge information have been treated as missing.

- In the last fortnight (11 August to 24 August 2025), there were fewer COVID-19 cases occupying intensive care beds across Australia than in the previous fortnight (Figure 18).
- In the last fortnight, there were fewer intensive care staff unavailable to work due to illness across Australia than in the previous fortnight (Figure 18).

Figure 18: Average number of COVID-19 cases in intensive care and the average number of intensive care staff unavailable to work due to illness by week of report*, Australia, 1 January to 24 August 2025




Source: Critical Health Resource Information System (CHRIS)

* Average number of ventilated and non-ventilated COVID-19 cases in intensive care includes only active COVID-19 cases (those in isolation) and does not include cleared COVID-19 cases.

† Intensive care staff include both medical and nursing staff. Staff unavailability will be underestimated in NSW as most public hospitals in NSW do not report staff unavailability.

- In the last fortnight, the number of COVID-19 cases occupying intensive care beds decreased in NSW, Qld, Tas, Vic and WA, and remained stable in the other jurisdictions compared with the previous fortnight (Figure 19).
- In the last fortnight, the number of intensive care staff unavailable to work due to illness increased in the ACT, the NT, SA and Tas but decreased or remained stable in the remaining jurisdictions compared with the previous fortnight (Figure 19).

Figure 19: Average number of COVID-19 cases in intensive care and the average number of intensive care staff unavailable to work due to illness by jurisdiction and week of report*†‡, Australia, 1 January to 24 August 2025



Source: Critical Health Resource Information System (CHRIS)

* Axis varies between jurisdictions.

† Average number of ventilated and non-ventilated COVID-19 cases in intensive care includes only active COVID-19 cases (those in isolation) and does not include cleared COVID-19 cases.


‡ Intensive care staff include both medical and nursing staff. Staff unavailability will be underestimated in NSW as most public hospitals in NSW do not report staff unavailability.

Mortality surveillance

Death registrations can provide information on the scale and severity of disease associated with acute respiratory infections. For more information on death registrations including completeness, timeliness, and definitions of deaths involving (both *due to* and *with*), *due to* and *with* acute respiratory infections, refer to the [Technical Supplement](#).

- COVID-19 has been the leading cause of acute respiratory infection mortality across 2023–2025.
- Since the end of 2021, a pattern has been observed for COVID-19 where there are two peaks of mortality during the year - one occurring between November and January and the other occurring between May and August. While there was still an increase in deaths involving COVID-19 occurring between November 2024 and January 2025, the number of deaths occurring during this period was much lower than other years (Figure 20a/b).
- The number of deaths *due to* COVID-19 increased in June 2025 following the relatively low number of deaths *due to* COVID-19 each month between February and May. Deaths remain at lower levels than the same period in 2024 and 2023 (Figure 20a).

Figure 20a: Provisional numbers of deaths *due to* an acute respiratory infection*† by month, year, and disease, Australia, 1 January 2023 to 31 July 2025




Source: Australian Bureau of Statistics, [Deaths due to acute respiratory infections in Australia](#), released 29 August 2025.

* Axis varies between acute respiratory infections.

† Data is provisional and subject to change. It can take several weeks for death registrations to be reported, processed, coded, validated, and tabulated. Therefore, the data shown here may be incomplete. Data for some months were not published by the ABS due to small counts, and therefore not reported here. Data includes all deaths (both doctor and coroner certified) that occurred and were registered by 31 July 2025.

- Deaths *due to* influenza increased in June 2025 and the number of deaths that occurred in July is likely to increase further (Figure 20a).
- There were 373 deaths *due to* influenza in the first six months of 2025 (Figure 20a). Although higher than the number of deaths *due to* influenza in the same period in 2023 (174) or 2024 (253), it remains lower than the 436 deaths recorded in the first six months of 2019, which was a recent high mortality year for influenza.
- There were 81 deaths *due to* RSV in the first six months of 2025; however, for privacy reasons six deaths *due to* RSV are not published in Figure 20a.
- The mortality rate for deaths *due to* COVID-19 or influenza for Aboriginal and Torres Strait Islander people was higher than for non-Indigenous people across each year in 2022–2024.
- Deaths *with* COVID-19 increased in June 2025 but remained lower than in 2023 or 2024 (Figure 20b).
- Deaths *with* influenza in May and June 2025 remained below the number recorded in April 2025 (Figure 20b).
- Deaths *with* RSV have increased each month from February to June 2025, similar to the increasing trend during the same period in 2023 and 2024 (Figure 20b).
- All three of these acute respiratory infections are more likely to cause death in older age groups than younger age groups.

Figure 20b: Provisional numbers of deaths *with* an acute respiratory infection*† by month, year, and disease, Australia, 1 January 2023 to 31 July 2025



Source: Australian Bureau of Statistics, [Deaths due to acute respiratory infections in Australia](#), released 29 August 2025.

* Axis varies between acute respiratory infections.


† Data is provisional and subject to change. It can take several weeks for death registrations to be reported, processed, coded, validated, and tabulated. Therefore, the data shown here may be incomplete. Data for some months were not published by the ABS due to small counts, and therefore not reported here. Data includes all deaths (both doctor and coroner certified) that occurred and were registered by 31 July 2025.

Laboratory surveillance

Sentinel laboratory surveillance monitors and characterises respiratory pathogens to provide information on what pathogens are circulating, potential changes in the pathogens that might affect their infectiousness, severity, ability to evade vaccine and/or infection-acquired immunity, or resistance to antivirals.

- In the last fortnight (11 August to 24 August 2025), SARS-CoV-2 test positivity decreased to 2.3% (333/14,704), influenza test positivity decreased to 12.2% (2,160/17,767), and RSV test positivity increased slightly to 6.0% (875/14,704) (Figure 21).

Figure 21: Number of tests positive (bars) and test positivity (line) for SARS-CoV-2, influenza or RSV of those specimens tested by sentinel laboratories by week of report[†], Australia, 1 January to 24 August 2025




Source: Sentinel laboratories, including National Influenza Centres

* Number of specimens tested excludes data from WA as testing denominator data are different for the three pathogens in Western Australia.

† A small minority of total samples from Victoria are tested only by respiratory panel (influenza, parainfluenza, adenovirus, human metapneumovirus, seasonal coronaviruses, RSV, and some picornaviruses) but not for SARS-CoV-2. These minority samples include only forensic materials; all other samples are tested by respiratory panel and SARS-CoV-2 assay.

- Please note, AusTrakka SARS-CoV-2 sequencing data are updated each month, as such SARS-CoV-2 sequence data presented here have not been updated since the previous report.
- There were 198 SARS-CoV-2 sequences uploaded to AusTrakka with dates of collection in the last 28 days (14 July to 10 August 2025). These sequences were from NSW, Qld, SA, Tas and WA, with the most recent collection date 2 August 2025.
- Most sequences were assigned to the BA.2.86 sub-lineage within B.1.1.529 (Omicron), or recombinants consisting of one or more Omicron sub-lineages (Figure 22a/b). In the last 28 days:
 - 17.7% (35/198) of sequences were from the sub-sub-lineages JN.1 (BA.2.86.1.1)
 - 81.8% (162/198) of sequences were recombinant or recombinant sub-lineages, including NB.1.8.1
 - there was one BA.3.2 sequence identified in the past 28 days
 - there were no BA.1, BA.4, BA.5 or other BA.2 sub-sub-lineage sequences.
- NB.1.8.1 is the dominant sub-lineage in the last 28 days, accounting for 72.2% (143/198) of sequences (Figure 22a).
- The World Health Organization (WHO) have identified certain sub-sub-lineages and recombinants as variants under monitoring (VUM) or variants of interest (VOI) because of their epidemiological, pathological, or immunological features of concern. A select number are highlighted below due to their relevance in the Australian context. There are:
 - 160 XFG sequences in AusTrakka, with 17 collected in the last 28 days
 - 1,394 NB.1.8.1 sequences in AusTrakka, with 143 collected in the last 28 days
 - 752 LP.8.1 sequences in AusTrakka, with 13 collected in the last 28 days
 - 3,086 KP.3.1.1 sequences in AusTrakka, with 13 sequences collected in the last 28 days
 - 3,462 XEC sequences in AusTrakka, with 1 sequence collected in last 28 days.

Figure 22a: SARS-CoV-2 Omicron sub-lineage* sequences by sample collection date, showing the proportions of sequences per week^{†‡}, Australia, 1 January to 10 August 2025




Source: AusTrakka

* Some sub-sublineages are shown alongside their parent lineage, but not included in the parent lineage totals. For instance, KP.2 and KP.3 are sub-sub lineages of JN.1, so the total of JN.1 sequences will be higher than shown in the corresponding colour alone, and should include the KP.2 and KP.3 totals.

† Sequences in AusTrakka aggregated by week and reported based on date of sample collection, not date of sequencing.

‡ Proportions in Figure 22a may not be representative when sequence numbers are small; refer to Figure 22b. Data for earlier weeks may change between reporting periods as sequences with older collection dates are uploaded. These numbers are not equivalent to number of cases, as there are many cases which may not be sequenced. Non-VOI and non-VUM Omicron sub-lineages have been collapsed into parent lineages BA.1, BA.2, BA.3, BA.4 and BA.5.

Figure 22b: SARS-CoV-2 Omicron sub-lineage* sequences by sample collection date, showing the count of sequences per week^{†‡}, Australia, 1 January to 10 August 2025



Source: AusTrakka

* Some sub-sublineages are shown alongside their parent lineage, but not included in the parent lineage totals. For instance, KP.2 and KP.3 are sub-sublineages of JN.1, so the total of JN.1 sequences will be higher than shown in the corresponding colour alone, and should include the KP.2 and KP.3 totals.

† Sequences in AusTrakka aggregated by week and reported based on date of sample collection, not date of sequencing.

‡ Data for earlier weeks may change between reporting periods as sequences with older collection dates are uploaded. These numbers are not equivalent to number of cases, as there are many cases which may not be sequenced. Non-VOI and non-VUM Omicron sub-lineages have been collapsed into parent lineages BA.1, BA.2, BA.3, BA.4 and BA.5.

- In the year to date, the WHO Collaborating Centre for Reference and Research on Influenza has antigenically characterised 3,370 influenza viruses from Australia (Table 4), of which:
 - 73.0% (2,459/3,370) have been influenza A(H1N1)
 - 8.7% (294/3,370) have been influenza A(H3N2)
 - 18.3% (617/3,370) have been influenza B/Victoria.
- In the year to date, there have been no influenza B/Yamagata viruses characterised (Table 4). The last influenza B/Yamagata virus characterised in Australia was in a sample from 2020.
- Of the influenza A(H1N1) samples tested for neuraminidase inhibitor resistance, 1.3% (14/1,061) demonstrated highly reduced inhibition to Oseltamivir. None of the influenza A(H3N2) samples tested for neuraminidase inhibitor resistance demonstrated highly reduced inhibition to Oseltamivir.
- None of the samples tested demonstrated highly reduced inhibition to Zanamivir.

Table 4: Australian influenza viruses typed by haemagglutination inhibition assay and jurisdiction*[†], 1 January to 24 August 2025

Strain	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
A(H1N1)	267	301	576	124	74	396	657	64	2,459
A(H3N2)	15	27	121	20	4	27	73	7	294
B/Victoria lineage	91	126	71	25	21	39	209	35	617
B/Yamagata lineage	0	0	0	0	0	0	0	0	0
Total	373	454	768	169	99	462	939	106	3,370

Source: World Health Organization (WHO) Collaborating Centre for Reference and Research on Influenza

*Viruses tested by the WHO Collaborating Centre for Reference and Research on Influenza are not necessarily a random sample of all those in the community and early-year data may be based on limited samples received. There may be up to a month delay on reporting of samples.

† Jurisdiction indicates the residential location for the individual tested, not the submitting laboratory.

Vaccine coverage, effectiveness and match

Vaccine coverage, effectiveness and match for acute respiratory infections are monitored from several data sources in Australia. Refer to the [Technical Supplement](#) for more information.

Vaccine coverage

- In Australia, regular COVID-19 vaccination is the best way to stay protected against severe illness, hospitalisation and death from COVID-19. Most adults should get vaccinated annually and adults aged 75 years and over should get vaccinated every six months.
 - More information on COVID-19 vaccines in Australia is available via the [department's COVID-19 webpages](#) or from the [National Centre for Immunisation Research and Surveillance \(NCIRS\)](#).
- Nationally, 8.9% of adults (aged 18 years and over) have received a COVID-19 vaccine in the last six months (Table 5).
- Nationally, fewer adults have received a COVID-19 vaccine in the last 12 months (11.3%; Table 5), compared to the 12 months prior (14.3% from 21 August 2023 to 18 August 2024).
- In the last 12 months, vaccine coverage decreased in all age groups, with the largest decrease seen in 65–74 years age group (from 34.7% in the 12 months prior to 27.5% in the last 12 months).
- There is substantial variation in COVID-19 vaccine coverage across age groups, ranging from 4.8% in adults aged 18–64 years to 43.4% in adults aged 75 years and over. COVID-19 vaccine coverage increases with increasing age (Table 5).
- There is some variation in vaccine coverage across jurisdictions, ranging from 4.3% in the NT to 19.3% in Tas (Table 5).

Table 5: COVID-19 vaccine coverage*†‡ by age group and jurisdiction, Australia, 19 August 2024 to 24 August 2025

Age group	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Last 12 months (19 August 2024 to 24 August 2025)									
18–64 years	10.6	4.1	2.2	4.6	4.7	8.7	5.2	4.8	4.8
65–74 years	48.6	25.4	15.3	26.4	28.3	40.6	27.9	28.0	27.5
≥ 75 years	67.7	41.1	28.1	42.5	43.2	58.4	42.6	45.2	43.4
All ages (18 years and over)	18.7	10.5	4.3	10.9	12.3	19.3	11.4	11.1	11.3
Last 6 months (24 February 2025 to 24 August 2025)									
18–64 years	8.3	3.1	1.5	3.5	3.6	6.8	4.0	3.9	3.7
65–74 years	39.5	20.6	12.0	20.8	23.1	32.9	22.7	23.2	22.2
≥ 75 years	54.9	32.6	22.5	33.0	34.6	46.4	33.7	36.4	34.3
All ages (18 years and over)	14.9	8.2	3.2	8.5	9.9	15.4	9.0	9.1	8.9

Source: Australian Immunisation Register (AIR) as at 25 August 2025

* COVID-19 vaccine coverage among the general population uses the most recently available Australian Bureau of Statistics Estimated Resident Population (ERP) as denominator for population data. Age in years is calculated as at the reporting week.

† COVID-19 vaccine coverage is influenced by changes in COVID-19 vaccine recommendations and eligibility criteria. For this reason, caution should be used when comparing coverage rates in the current 12 month period to previous 12 month periods. Coverage data in these tables may differ slightly from coverage estimates in other reports due to differences in calculation methodologies and/or different data download dates.

‡ Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Population denominator data used to calculate COVID-19 vaccine coverage are based on an individual's residential address. Total rows will include individuals where jurisdiction was missing.

- Nationally, 3.7% of Aboriginal and Torres Strait Islander adults (aged 18 years or over) have received a COVID-19 vaccine in the last six months (Table 6).
- Nationally, fewer Aboriginal and Torres Strait Islander adults have received a COVID-19 vaccine in the last 12 months (4.7%; Table 6), compared to the 12 months prior (6.7% from 21 August 2023 to 18 August 2024).
- In the last 12 months, vaccine coverage decreased in all age groups of Aboriginal and Torres Strait Islander people, with the largest decrease seen in the 75 years and over age group (from 34.3% in the 12 months prior to 27.3% in the last 12 months).
- Among Aboriginal and Torres Strait Islander people there is substantial variation in COVID-19 vaccine coverage across age groups, ranging from 2.7% in adults aged 18–64 years to 27.3% in adults aged 75 years and over. COVID-19 vaccine coverage increases with increasing age (Table 6).
- Among Aboriginal and Torres Strait Islander people, there is slight variation in vaccine coverage across jurisdictions, ranging from 2.6% in the NT to 9.7% in the ACT (Table 6).

Table 6: COVID-19 vaccine coverage*†‡ among Aboriginal and Torres Strait Islander populations by age group and jurisdiction, Australia, 19 August 2024 to 24 August 2025

Age group	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Last 12 months (19 August 2024 to 24 August 2025)									
18–64 years	6.4	2.7	1.9	2.6	2.7	5.5	4.0	2.2	2.7
65–74 years	35.7	19.0	8.6	17.6	17.5	32.7	20.7	16.2	18.0
≥ 75 years	51.1	29.4	13.3	25.7	29.3	43.6	32.2	26.2	27.3
All ages (18 years and over)	9.7	5.1	2.6	4.4	4.8	9.7	6.6	3.7	4.7
Last 6 months (24 February 2025 to 24 August 2025)									
18–64 years	5.1	2.1	1.2	2.0	2.1	4.3	3.0	1.8	2.1
65–74 years	27.7	15.4	6.5	14.0	14.3	27.0	16.9	13.4	14.5
≥ 75 years	40.7	23.2	11.2	20.2	22.9	33.8	25.4	21.5	21.6
All ages (18 years and over)	7.7	4.1	1.8	3.4	3.8	7.7	5.1	3.1	3.7

Source: Australian Immunisation Register (AIR) as at 25 August 2025

* COVID-19 vaccine coverage among Aboriginal and Torres Strait Islander populations is based on the AIR population as known at the reporting week. Age in years is calculated as at the reporting week.

† COVID-19 vaccine coverage is influenced by changes in COVID-19 vaccine recommendations and eligibility criteria. For this reason, caution should be used when comparing coverage rates in the 12 month period to previous 12 month periods. Coverage data in these tables may differ slightly from coverage estimates in other reports due to differences in calculation methodologies and/or different data download dates.

‡ Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Population denominator data used to calculate COVID-19 vaccine coverage are based on an individual's residential address. Total rows will include individuals where jurisdiction was missing.

- Influenza virus strains change year to year, so annual vaccination before the peak of the influenza season provides Australians with the best protection against influenza and its complications. The seasonal influenza vaccine is recommended for everyone aged six months and over.
 - More information on influenza vaccines in Australia is available via the [department's influenza vaccine webpages](#) or from [NCIRS](#).
- Nationally, influenza vaccine coverage is 30.2% for 2025 so far (Table 7); however, remains lower than coverage at the same time in 2023 and 2022.
- There is substantial variation in influenza vaccine coverage across age groups, ranging from 14.6% in children aged 5–14 years to 60.9% in adults aged 65 years and over (Table 7). The current trend should be interpreted with care as people aged 5–64 years are generally not eligible for free seasonal influenza vaccine under the National Immunisation Program.
- There is some variation in influenza vaccine coverage across jurisdictions, ranging from 23.8% in the NT to 39.9% in the ACT (Table 7).
- Among Aboriginal and Torres Strait Islander populations, there is substantial variation in influenza vaccine coverage across age groups, ranging from 11.7% in children aged 5–14 years to 60.4% in adults aged 65 years and over (Table 7).
- Among Aboriginal and Torres Strait Islander populations, there is some variation in influenza vaccine coverage across jurisdictions, ranging from 19% in WA to 30.5% in the NT (Table 7).

Table 7: Influenza vaccine coverage*†‡ by age group and jurisdiction, Australia, 1 March to 24 August 2025

	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Age groups									
6 months to <5 years	51.9	27.7	36.5	22.1	31.1	33.0	33.9	25.9	28.8
5–14 years	24.3	13.4	12.5	13.1	15.4	14.9	16.1	15.3	14.6
15–49 years	32.4	19.3	21.3	18.1	23.4	23.3	23.5	18.7	20.7
50–64 years	44.1	29.9	25.4	31.2	35.9	39.0	34.1	30.8	32.1
≥ 65 years	65.1	58.5	36.8	60.9	67.0	68.3	62.2	60.4	60.9
All ages (6 months and over)	39.9	28.8	23.8	28.1	34.7	36.2	32.3	28.3	30.2
Aboriginal and Torres Strait Islander populations									
6 months to <5 years	32.7	18.5	35.3	14.7	18.2	23.1	22.8	18.2	18.9
5–14 years	16.7	10.9	19.8	10.3	12.1	12.5	12.3	11.2	11.7
15–49 years	25.7	16.7	28.6	15.4	19.1	19.1	19.1	15.3	17.7
50–64 years	45.5	36.1	42.2	34.4	37.8	45.1	36.3	32.0	36.2
≥ 65 years	68.9	62.6	46.1	60.2	63.5	69.7	63.6	53.9	60.4
All ages (6 months and over)	29.7	21.6	30.5	19.4	23.1	26.0	24.2	19.0	21.9

Source: Australian Immunisation Register (AIR) as at 25 August 2025

* Influenza vaccine coverage uses the AIR population as the denominator. Coverage data in these tables may differ slightly from coverage estimates in other reports due to differences in calculation methodologies and/or different data download dates.

† Age is calculated based on the person's age as at 1 July of the reporting year.

‡ From the report ending 13 July 2025, jurisdiction is based on the person's address on the AIR rather than an individual's residential address as recorded on Medicare. Total rows will include individuals where jurisdiction was missing. In addition, to align with departmental reporting methodologies, both the numerator (number of persons vaccinated) and denominator (AIR population) for influenza vaccine coverage only consider person records with a Personal Identification Number that was able to be matched to Medicare. Person records with a Synthetic Identification Number are now excluded from both numerator and denominator. For these reasons, influenza vaccine coverage metrics in previous Australian Respiratory Surveillance Reports and coverage metrics from the report ending 13 July 2025 moving forward should be interpreted with care.

- Infants can be protected against severe illness from RSV through the vaccination of pregnant people or the direct administration of monoclonal antibodies like nirsevimab. These are part of the National RSV Maternal and Infant Protection Program which launched on 3 February 2025 and includes both the National Immunisation Program funded Abrysvo vaccine and jurisdictional nirsevimab programs.
 - More information on RSV immunisation in Australia is available via the [department's RSV vaccine webpages](#) or from [NCIRS](#).
- Since the commencement of the National RSV Mother and Infant Protection Program, 109,939 Abrysvo doses have been administered to pregnant people nationally (Table 8).
- While high maternal vaccine uptake is a positive indicator of maternal program success, it may result in lower nirsevimab uptake rates in infants. This is because maternal antibodies passed to the infant can provide protection against RSV, potentially reducing the need for infant immunisation.

Table 8: Number of doses of Abrysvo administered to pregnant people by jurisdiction*, Australia, 3 February to 24 August 2025

	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Age group									
15–24 years	176	2,793	196	2,421	607	297	1,593	1,063	9,146
25–39 years	2,657	30,008	875	16,929	6,606	2,014	26,042	9,729	94,860
40–54 years	174	1,968	42	891	374	92	1,791	601	5,933
Total (15–54 years)	3,007	34,769	1,113	20,241	7,587	2,403	29,426	11,393	109,939

Source: Australian Immunisation Register (AIR) as at 25 August 2025

* Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Total rows will include individuals where jurisdiction was missing.

- In the last six months, 21.0% of infants (aged < 8 months) have received nirsevimab (Table 9).
- There is substantial variation in nirsevimab uptake in infants across jurisdictions, ranging from 13.2% in NSW to 39.8% in WA (Table 9).
- The current trend is likely due to variation in the seasonality and eligibility criteria between state and territory programs, as well as the presence of previous nirsevimab programs. Some state and territory programs are seasonal (from 1 April to 30 September), whereas others are year-round. In states with seasonal programs (SA, Tas, Vic, and parts of WA), uptake may appear disproportionately lower at this time of the year. In addition, Qld and WA had nirsevimab programs in 2024, which may contribute to higher nirsevimab uptake in 2025 in these states.

Table 9: Nirsevimab (Beyfortus) uptake in the last six months*† by age group and jurisdiction, Australia, 24 February to 24 August 2025

	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Age group									
Infants (aged < 8 months)	16.6	13.2	21.9	18.9	31.3	34.8	21.3	39.8	21.0
Young children (aged ≥ 8 to 24 months)	1.1	0.4	0.4	0.2	1.1	1.5	1.0	1.3	0.7

Source: Australian Immunisation Register (AIR) as at 25 August 2025

* Reporting of RSV monoclonal antibodies to the AIR is not compulsory; therefore, uptake is likely to be underestimated. Uptake data in these tables may differ slightly from estimates in other reports due to differences in calculation methodologies and/or different data download dates.

† For infants and young children vaccinated, age in months is calculate as months between the immunisation encounter and date of birth rounded down as at the reporting date. For the infant and young children population, age in months is calculated as months between the AIR data extract date and date of birth rounded down as at the reporting date.

‡ Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Total rows will include individuals where jurisdiction was missing. Population denominator data used to calculate nirsevimab uptake are based on an individual's residential address as recorded on Medicare.

Vaccine effectiveness

- It is too early to assess vaccine effectiveness for the 2025 influenza season.

Vaccine match

- Refer to the [Technical Supplement](#) for information on the 2025 southern hemisphere influenza vaccines composition.
- In the year to date, 99.3% (2,442/2,459) of influenza A(H1N1) isolates, 99.3% (292/294) of influenza A(H3N2) isolates and 98.9% (610/617) of influenza B/Victoria lineage isolates characterised have been antigenically similar to the corresponding 2025 vaccine components.