s 22

From:	s 47F
Sent:	Friday, 26 July 2019 10:20 PM
То:	s 22
Subject:	oncotype DX [SEC=No Protective Marking]
Attachments:	cost effectiveness of chemo.pdf



KR s 47F Table 11 Summary of disaggreg	ated incremental cost	and effectiveness in C	T sparing only
Parameter	Oncotype DX	Usual care	Incremental
Disaggregated costs			
Oncotype DX test costs	\$5,085.00	\$0.00	\$5,085.00
Chemotherapy	\$1,253.65	\$3,116.03	-\$1,862.38
Hormone therapy	\$3,160.85	\$3,160.85	\$0.00
Recurrent disease	\$5,791.22	\$5,791.22	\$0.00
Total	\$15,290.72	\$12,068.10	\$3,222.62
Disaggregated outcomes (discounted with h	alf cycle correction)		
Life years	13.6530	13.6530	0
Disease free	13.4577	13.4577	0
Post recurrence	0.1953	0.1953	0
QALY	13.4621	13.4575	<mark>0.0045</mark>
Disease free	13.3066	13.3021	0.0045
Post recurrence	0.1554	0.1554	0
		\$ per life year gained	\$NA
		\$ per QALY gained	\$711,529

s 47F

CRICOS 00026A This email plus any attachments to it are confidential. Any unauthorised use is strictly proh bited. If you receive this email in error, please delete it and any attachments.

Please think of our environment and only print this e-mail if necessary

HISOCUMERED ARTING OF HEALTH



The cost-effectiveness of adjuvant chemotherapy for early breast cancer: A comparison of no chemotherapy and first, second, and third generation regimens for patients with differing prognoses $\stackrel{\sim}{\sim}$

H.E. Campbell ^{a,*}, D. Epstein ^b, D. Bloomfield ^c, S. Griffin ^b, A. Manca ^b, J. Yarnold ^d, J. Bliss ^e, L. Johnson ^e, H. Earl ^f, C. Poole ^g, L. Hiller ^h, J. Dunn ^h, P. Hopwood ^e, P. Barrett-Lee ⁱ, P. Ellis ^j, D. Cameron ^k, A.L. Harris ¹, A.M. Gray ^a, M.J. Sculpher ^b

^a Health Economics Research Centre, University of Oxford, Headington, Oxford, OX3 7LF, United Kingdom

^b Centre for Health Economics, University of York, Heslington, York, YO10 5DD, United Kingdom

^c Sussex Cancer Centre, Brighton & Sussex University Hospitals, Brighton, BN2 5BE, United Kingdom

^d Royal Marsden NHS Trust, London and Sutton, United Kingdom

^e ICR CTSU, Section of Clinical Trials, The Institute of Cancer Research, Sutton, Surrey, SM2 5NG, United Kingdom

^f University of Cambridge, Department of Oncology, NIHR Cambridge Biomedical Research Centre, Cambridge, CB2 0QQ, United Kingdom

^g University of Warwick, Warwick, Coventry, CV4 7AL, United Kingdom

h Warwick Medical School Clinical Trials Unit, University of Warwick, Coventry, CV4 7AL, United Kingdom

ⁱ Velindre Cancer Centre, Velindre Hospital, Cardiff, CF14 2TL, United Kingdom

^j Guys and St Thomas NHS Foundation Trust, Cancer Management Office, Guys Hospital, London, SE1 9RT, United Kingdom

^k Edinburgh Cancer Research Centre, Western General Hospital, Edinburgh, EH4 2XR, United Kingdom

¹ Cancer Research UK Medical Oncology Department, Churchill Hospital, Headington, Oxford, OX3 7LJ, United Kingdom

ARTICLE INFO

Article history: Available online 7 July 2011,⁄

Keywords:

Cost effectiveness Decision analytic modelling Early breast cancer Chemotherapy

ABSTRACT

Background: The risk of recurrence following surgery in women with early breast cancer varies, depending upon prognostic factors. Adjuvant chemotherapy reduces this risk; how ever, increasingly effective regimens are associated with higher costs and toxicity profiles, making it likely that different regimens may be cost effective for women with differing prognoses. To investigate this we performed a cost effectiveness analysis of four treatment strategies: (1) no chemotherapy, (2) chemotherapy using cyclophosphamide, methotrexate, and fluorouracil (CMF) (a first generation regimen), (3) chemotherapy using Epirubicin CMF (E CMF) or fluorouracil, epirubicin, and cyclophosphamide (FEC60) (a second generation regimens), and (4) chemotherapy with FEC60 followed by docetaxel (FEC D) (a third gener ation regimen). These adjuvant chemotherapy regimens were used in three large UK led randomised controlled trials (RCTs).

Methods: A Markov model was used to simulate the natural progression of early breast can cer and the impact of chemotherapy on modifying this process. The probability of a first recurrent event within the model was estimated for women with different prognostic risk profiles using a parametric regression based survival model incorporating established

E mail address: helen.campbell@dph.ox.ac.uk (H.E. Campbell).

0959 8049/\$ see front matter © 2011 Elsevier Ltd. All rights reserved. doi:10.1016/j.ejca.2011.06.019

^{*} On behalf of the Centre for Health Economics, York, the ABC, NEAT and TACT Trial groups and the Health Economics Research Centre, University of Oxford.

^{*} Corresponding author: Address: Health Economics Research Centre, Department of Public Health, University of Oxford, Old Road Campus, Headington, Oxford, OX3 7LF, United Kingdom. Tel.: +44 1865 289274; fax: +44 01865 289271.

prognostic factors. Other probabilities, treatment effects, costs and quality of life weights were estimated primarily using data from the three UK led RCTs, a meta analysis of all rel evant RCTs, and other published literature. The model predicted the lifetime costs, quality adjusted life years (QALYs) and cost effectiveness of the four strategies for women with dif fering prognoses. Sensitivity analyses investigated the impact of uncertain parameters and model assumptions.

Findings: For women with an average to high risk of recurrence (based upon prognostic fac tors and any other adjuvant therapies received), FEC D appeared most cost effective assuming a threshold of £20,000 per QALY for the National Health Service (NHS). For youn ger low risk women, E CMF/FEC60 tended to be the optimal strategy and, for some older low risk women, the model suggested a policy of no chemotherapy was cost effective. For no patient group was CMF chemotherapy the preferred option. Sensitivity analyses demonstrated cost effectiveness results to be particularly sensitive to the treatment effect estimate for FEC D and the future price of docetaxel.

Interpretation: To our knowledge, this analysis is the first cost effectiveness comparison of no chemotherapy, and first, second, and third generation adjuvant chemotherapy regimens for early breast cancer patients with differing prognoses. The results demonstrate the potential for different treatment strategies to be cost effective for different types of patients. These findings may prove useful for policy makers attempting to formulate cost effective treatment guidelines in the field of early breast cancer.

💑 2011 Elsevier Ltd. All rights reserved.

1. Introduction

For women with early breast cancer, surgery to remove the pri mary tumour is often followed by adjuvant chemotherapy, a combination of cytotoxic drugs given to reduce the risk of dis ease recurrence. Recent decades have seen major advances in chemotherapy treatment as new and potentially more effective classes of drugs and modes of administration are developed. Randomised controlled trials (RCTs) have evaluated first gener ation alkylating based chemotherapy regimens, second gener ation anthracycline based regimens and third generation taxane based regimens.^{1–5}

Health economic evaluation has been used to assess the cost effectiveness of adjuvant chemotherapy regimens; most recently, those containing the taxane agents docetaxel and paclitaxel.^{6–9} However, these evaluations, and the RCTs provid ing the corresponding clinical data, conducted pair wise com parisons in which the new generation therapy was assessed only against the previous generation's therapy. To our knowl edge, no clinical or economic assessment has to date simulta neously compared all potential adjuvant chemotherapy treatment strategies; that is, first, second, and third generation regimens, as well as a policy of using no adjuvant chemother apy. A simultaneous comparison of this kind would prove infor mative for several reasons. Firstly, there has been considerable recent research on the effectiveness of all adjuvant chemother apies, including meta analyses conducted by the Early Breast Cancer Trialists' Collaborative Group (EBCTCG) and systematic reviews by the Cochrane Breast Cancer Group.^{1,10} This evidence should inform and update economic analysis and policy. Sec ondly, there are factors relating to the progression and treat ment of early breast cancer that mean different chemotherapy regimens may be cost effective for different pa tient groups. The risk of recurrence, for example, can vary sub stantially between women depending upon prognostic factors

including age, axillary lymph node status, and tumour grade, size, and oestrogen receptor (ER) status.^{11–13} Also, the propor tional reduction in the risk of recurrence with chemotherapy is known to vary by age, and women with ER positive tumours now routinely receive adjuvant anti hormone therapy, which also reduces the risk of recurrence, but independently of che motherapy.¹ Finally, advances in treatment effectiveness come at a cost, as new chemotherapy regimens are usually more expensive than older regimens and potentially more toxic, resulting in an increase in adverse events and a detrimental im pact on patient health related quality of life (HRQoL). Given all of these factors, it would appear important to review each of the relevant treatment options for all types of patient.

This paper describes a cost effectiveness analysis in which the lifetime costs and quality adjusted life years (QALYs) (life years adjusted for HRQoL) of adjuvant chemotherapy treat ment options for women with early breast cancer are estimated and compared. Reflecting the full range of management op tions which have been used in the adjuvant setting in United Kingdom trials over recent years, the strategies considered are:

- I. no chemotherapy,
- II. first generation chemotherapy with CMF (six cycles of cyclophosphamide, methotrexate, fluorouracil),
- III. second generation chemotherapy with FEC60 (eight cycles of fluorouracil, epirubicin, cyclophosphamide) or E CMF (four cycles of epirubicin followed by four cycles of CMF), and
- IV. third generation chemotherapy with FEC D (four cycles of FEC60 followed by four cycles of docetaxel).

These chemotherapy regimens are defined in accordance with the protocols of three pragmatic UK led RCTs the Adju vant Breast Cancer (ABC) Trial which began in 1992 and com pared CMF with no chemotherapy, the National Epirubicin Adjuvant Trial (NEAT) which began in 1996 and compared E CMF with CMF, and the Taxotere as Adjuvant Chemotherapy Trial (TACT) which began in 2001 and compared FEC D with FEC60 or E CMF.^{3–5} Docetaxel is not currently licensed in the UK for use as a single agent following FEC. This regimen is, however, advocated by the majority of breast cancer oncolo gists in the UK and indeed worldwide, because when used on its own, docetaxel carries a lower risk of adverse events than if used simultaneously with other agents.¹⁴ In line with the TACT trial protocol and indirect evidence from the trial re sults, our analysis assumes equivalence of FEC60 and E CMF in terms of effectiveness and toxicity.

2. Methods

2.1. Overview and model structure

The analysis followed the methodological guidelines from the National Institute of Health and Clinical Excellence (NICE).¹⁵ It was conducted from a UK National Health Service (NHS) per spective and the annual discount rate used for costs (ex pressed as 2009 Pound Sterling) and effects was 3.5%.¹⁶ Fig. 1 shows the structure of the economic model. Following local surgery, (a hypothetical cohort of) patients enter a short term decision tree (panel a) and are assigned to one of the



Panel A: decision tree for first 6 months of the analysis

*Models with this structure are attached to each of the four branches in the decision tree.

four treatment options being compared. During the first 6 months (the average duration of chemotherapy) they are as sumed to be free from breast cancer. Patients receiving che motherapy incur treatment costs and face the risk of regimen specific toxicity which may culminate in death, hos pitalisation and/or reduced HRQoL.

In month seven, patients enter the Markov model (panel b) which simulates the natural progression of the disease, and the impact of chemotherapy on modifying this process. The model comprises a set of mutually exclusive health states, which patients can move between at annual intervals accord ing to transition probabilities. Values are attached to each health state in the model to reflect the cost and HRQoL asso ciated with a year spent in that state. Patients then accrue costs and QALYs according to their health state in the model during each year.

Following chemotherapy, surviving patients enter the 'dis ease free' health state of the model where they remain until experiencing a first recurrent breast cancer event (which can be local, metastatic, or a contralateral/second primary breast tumour) or death from causes unrelated to breast can cer. Patients transiting to a recurrence health state remain there until they die from breast cancer or another cause. Although further recurrent events experienced prior to a breast cancer death (for example a metastatic recurrence fol lowing a local recurrence) are not explicitly modelled with additional health states (this would over complicate the mod el's structure), their consequences are still captured within the analysis through the cost and HRQoL values assigned to the existing recurrence health states. The model was devel oped and run in Excel 2007 (Microsoft Corporation, Redmond, WA, United States of America).

2.2. Populating the model

Parameters were estimated using individual patient data col lected as part of the ABC, NEAT, and TACT trials, and from administrative datasets, comprehensive searches of the pub lished literature, and expert clinical opinion. Abridged detail is given below, with more information available in the web appendices.

2.3. During chemotherapy

Based on individual patient data from the three trials, Table 1 presents the expected cost per patient of each chemotherapy regimen and of breast cancer related hospital admissions dur ing chemotherapy. Informed by data collected during the ABC trial, women entering the model were assumed to have HRQoL on average equivalent to women in the age matched UK general population. This is based on the EQ 5D instru ment, a generic measure of HROoL consisting of five dimen sions (mobility, self care, usual activities, pain/discomfort, and anxiety/depression) on each of which the respondent indicates whether they have no problems, some problems or severe problems.¹⁷ These responses are translated onto a val uation scale between 0 (equivalent to death) and 1 (equivalent to full health) based on the views of a sample of the UK pub lic.18 During chemotherapy, the decrements in HRQoL (again using the EQ 5D) were estimated for each regimen based on trial data (Table 1). Patients may die as a result of chemother apy related toxicity in the first 6 months. Across the three RCTs such deaths were rare and so a pooled probability of tox icity related death was calculated and applied to all chemo therapy patients (Table 1).

2.4. After chemotherapy

2.4.1. Transition probabilities

Table 2 gives the transition probabilities used in the Markov model.

The underlying annual probability of a first recurrent event was derived from a published and validated parametric regression based survival model (see web Appendix 1).¹⁹ Risk

Table 1 – Mean per patient cost and HRQoL values used to populate the decision tree. ^a				
Parameter	Value	Source		
Costs				
CMF chemotherapy	£3,029	ABC, ^{39–42}		
E CMF/FEC60 chemotherapy	£3,576	TACT, ^{39–42}		
FEC D chemotherapy	£6,936	TACT, ^{39–42}		
Breast cancer related hospital admissions during CMF	£90	NEAT, ⁴³		
Breast cancer related hospital admissions during E CMF/I	FEC60 £123	NEAT, ⁴³		
Breast cancer related hospital admissions during FEC D	£175	TACT, ⁴³		
HRQoL based on the EQ 5D health instrument	Age and sex adjusted	44		
Underlying HRQoL levels	United Kingdom			
	population norms			
Decrement in HRQoL from underlying levels in				
Patients not receiving chemotherapy	.003	ABC		
Patients receiving CMF	0.057	ABC		
Patients receiving E CMF/FEC60	0.067	NEAT, TACT		
Patients receiving FEC D	0.099	TACT		
Probability of death from chemotherapy related	0.00371	ABC, NEAT, TACT		
toxicity during 6 months of treatment				

^a Parameter estimates shown are based upon analyses of individual patient data collected during the three trials and are comprised of multiple components (costs) or are additive in nature (HRQoL decrements). Individual components and estimates of their associated uncertainty are given in web Appendix 1.

Table 2 – Parameter values used within the Markov model.

Parameter	Value	Source
Annual transition probabilities First recurrent event in the absence of hormone therapy and chemotherapy (predictions include radiotherapy) years 1 50	Variable dependent upon age, number of positive nodes, grade, size, and ER status of the primary tumour, and	19
First recurrent event is local First recurrent event is metastatic First recurrent event is a contralateral/second primary breast tumour	time since diagnosis 0.227 (461/2028) 0.705 (1430/2028) 0.068 (137/2028)	ABC, TACT, data from Churchill Hospital, Oxford
Breast cancer death following recurrence years 1 15 post recurrence	Variable dependent upon type of recurrence and time since	ABC, TACT, data from Churchill Hospital, Oxford
Breast cancer death following recurrence years 16 and onwards post recurrence	4.7 times age adjusted UK breast cancer mortality	20,22
Death from causes other than breast cancer years 1 50	Age and sex adjusted UK	21,22
Relative treatment effect estimates for recurrence Tamoxifen versus no hormone therapy (ER positive	0.59 (SE 0.03)	1
Aromatase inhibitor versus tamoxifen (ER positive	0.74 (SE 0.06) ^d	45
CMF versus no chemotherapy, women aged <50 years	0.59 (SE 0.04)	1
CMF versus no chemotherapy, women aged >50 years	0.81 (SE 0.03)	1
Anthracycline based chemotherapy versus CMF ^b years	0.89 (SE 0.03)	1
1 10 FEC D versus E CMF/FEC60 ^c years 1 10	0.90 (SE 0.05)	2,3
Mammogram	£46.37 (SF £9.27)	42,46
Outpatient clinic attendance	£99.00 (SE £19.80)	43
Local recurrence vear 1	£13,898 (SE £2,388) ^d	24,42
Local recurrence years 2 5	£1,235 (SE £428) ^d	24,42
Contralateral/second primary breast tumour year 1	£18,374 (SE £4,750) ^d	24,42
Contralateral/second primary breast tumour years 2 5	£586 (SE £101) ^d	24,42
Metastatic recurrence year 1	£10,482 (SE £1,135) ^d	24,42
Metastatic recurrence year 2	£8,609 (SE £1,354) ^a	24,42
Metastatic recurrence year 3	£4,811 (SE £1,506) ^a	24,42
Metastatic recurrence year 4	12,270 (SE 1880)	24,42
End of life care (3 months prior to breast cancer death)	$f_{4,038} (SF f_{454})^{d}$	24,42
HROOL	1,000 (01 1104)	
Underlying level of HRQoL	Population norms	44
Decrement in HRQoL with CMF year 1 ^e	0.057	ABC
Decrement in HRQoL with E CMF/FEC60 year 1 ^e	0.038	NEAT, TACT
Decrement in HRQoL with FEC D year 1 ^e	0.035	TACT
Decrement in HRQoL with local recurrence	0.108 (SE 0.04)	47-50
Decrement in HRQoL with metastatic recurrence	0.303 (SE 0.16)	47-50
breast tumour	0.108 (SE 0.04)	
HRQoL in final 3 months of life	0.159 (SE 0.04)	52,55-57

^a Treatment effect for aromatase inhibitor is for anastrazole and is taken from the ATAC trial.⁴⁵

^b Used in this analysis as effectiveness estimate for E CMF/FEC60.

^c The hazard ratio for FEC D versus E CMF/FEC60 is pooled from the TACT and PACS01 trials and is for disease free survival (recurrence plus death from any cause) rather than just recurrence. Given the small number of deaths from treatment related toxicity, and an expectation that deaths from other causes would be equally distributed across both arms of the TACT and PACS01 trials, we assume the ratio for DFS is a reasonable proxy for the ratio for first recurrence, the parameter used in the model.

^d Standard error inferred from 95% confidence intervals.

^e HRQoL decrements in the year following chemotherapy are additive in nature. Individual components and estimates of their associated uncertainty are given in web Appendix 1.

of recurrence was predicted on the basis of a patient's age, number of positive axillary lymph nodes, the grade, size, and ER status of the primary tumour, assuming radiotherapy is given. The predictions are time dependent, to reflect the increasing risk of recurrence in the first 2 years after breast cancer surgery and the decreasing risk thereafter. Following the prediction of a first recurrent event in a given year, pa tients are distributed across the Markov model's three recur rence health states according to the proportions shown in Table 2.

The annual risk of breast cancer death following each type of first recurrent event was estimated using a second para metric regression based survival model (see web Appendix 1). This model was estimated using patient data from a num ber of sources and is time dependent to allow for the increas ing mortality risk in the first 3 4 years after recurrence, and a declining risk thereafter. It was used to predict the probability of breast cancer death for the first 15 years after each type of recurrence. Thereafter the risk of death following breast can cer was based on the rate for the UK general population in flated by a factor of 4.7 as observed for long term breast cancer survivors (see Table 2).²⁰

Finally, deaths from causes other than breast cancer were accounted for in the model by applying age and sex adjusted UK life table data (excluding breast cancer deaths) to all non terminal health states.^{21,22}

2.4.2. Treatment effects

Table 2 shows the treatment effect estimates, in the form of hazard ratios, used to adjust the underlying annual risk of a recurrent event in the first 10 years in the model. As per

clinical practice (and independent of any chemotherapy mod elled) pre menopausal women with ER positive tumours were assumed to receive 5 years of tamoxifen and post meno pausal ER positive women 5 years with an aromatase inhibi tor (see Table 2). Estimates of the effectiveness of CMF and E CMF/FEC60 regimens were taken from the Early Breast Can cer Trialists' Collaborative Group (EBCTCG) meta analyses; for FEC D estimates were taken from a pooled analysis of data from the only two trials to have compared the sequential use of docetaxel with an anthracycline based comparator of a similar duration (TACT and PACS 01).^{1–3} The effect of each type of chemotherapy was on the risk of recurrence, and for E CMF/FEC60 and FEC D, was estimated indirectly by assum ing hazard ratios were additive on the log scale (see Fig. 2). In line with the published literature, the model employed sepa rate chemotherapy treatment effect estimates for women above and below the age of 50.¹

2.4.3. Costs

Patients who remained in remission or survived more than 5 years following a local or contralateral recurrence or second primary tumour were assumed to undergo mammographic surveillance and outpatient follow up according to NICE guidelines.²³ The unit costs applied to these contacts are shown in Table 2.

A published costing study was used as the source of an nual costs of treating each type of recurrence for 5 years and for an estimate of the cost of terminal care which was as signed to patients in the model 3 months prior to a breast cancer death (see Table 2).²⁴ Patients in that study are likely to be representative of patients in the ABC, NEAT and TACT

Direct evidence is available on the effectiveness of CMF (B) versus No chemotherapy (A), E-CMF/FEC60 (C) versus CMF (B), and FEC-D (D) versus E-CMF/FEC60 (C).



The effectiveness of C versus A can be estimated indirectly as follows:

$$\mathsf{LHR}_\mathsf{CA} = \mathsf{LHR}_\mathsf{BA} + \mathsf{LHR}_\mathsf{CB}$$

The effectiveness of D versus A can be estimated in the same way:

$$LHR_{DA} = LHR_{BA} + LHR_{CB} + LHR_{DC}$$

Where LHR = logged hazard ratio

Adapted from: Caldwell D. Introduction to indirect and mixed treatment comparisons (MTC). Multiple treatment comparisons workshop for NICE. September 1st 2006.³⁸

Fig. 2 – Indirect estimation of the effect of E-CMF/FEC60 and FEC-D upon the underlying risk of recurrence.³⁸

FOI 1513

trials, having been treated at a UK oncology department that recruited patients to all three of these studies. For patients with a local recurrence or a contralateral/second primary breast tumour and who died from breast cancer at the end of a year, metastatic disease was assumed to have been diag nosed half way through that year, and related treatment costs (and HRQoL decrements see below) applied.

2.4.4. HRQoL

Data collected from ABC, NEAT, and TACT suggested the neg ative impact of chemotherapy on HRQoL continues for up to a year following treatment in patients who remain recurrence free (see Table 2). HRQoL decrements applied to patients in the recurrence health states of the model and in the 3 months preceding a breast cancer death were informed by a review of published literature (see Table 2 and web Appendix 1).

2.5. Cost-effectiveness

The time horizon for the analysis, including the first 6 months in the decision tree, was 50.5 years. The discounted costs and QALYs accrued by patients during these years were summed to give an estimate of expected lifetime costs and effects with each treatment strategy. All strategies were then compared and those that were dominated (i.e. more costly but no more effective than another strategy or a combination of other strategies) were excluded. Remaining strategies were ranked in increasing order of effectiveness and incremental cost effectiveness ratios (ICERs) computed.²⁵ The ICER provides an estimate of the additional cost of generating one addi tional QALY, and is calculated as the difference in cost divided by the difference in QALYs between a treatment and its next best alternative. These ICERs were compared against the cost effectiveness threshold, which, in the case of NICE in the UK, lies between £20,000 and £30,000 per QALY and seeks to reflect the health foregone when new interventions are funded and displace existing services.¹⁵ The most effective strategy with an ICER below this threshold is considered most cost effective.

For every level of prognosis, the potential exists for a range of cost effectiveness results depending upon the age (which determines the magnitude of chemotherapy effectiveness) and ER status (which determines the use of anti hormone therapy) of the patient. For this reason a definitive 'base case' result is not presented. Rather, cost effectiveness is first estimated for a cohort of younger (age 40) ER negative women. The number of positive nodes (1), tumour grade (2), and tu mour size (3 cm) for this cohort were selected so that, to gether with age and ER status, they generated a baseline prognosis prediction equivalent to that of the 'average risk' patient in the ABC trial, where the probability of remaining recurrence free out to 10 years in the absence of chemother apy and anti hormone therapy (as predicted by the model on the basis of prognostic factors and assuming radiotherapy is given) was estimated to be around 44%. In a series of sub group analyses, the impact of increasing the age and chang ing the ER status of this 'reference case' cohort is explored.

2.6. Sub-group analyses

The age of the reference case cohort was increased from 40 to 60 to assess the impact of reduced chemotherapy effective ness in older women. In both of these age groups ER status was also switched from negative to positive, and the assump tion made was that younger women would receive 5 years of tamoxifen and older women 5 years of an aromatase inhibi tor. Additional analyses were conducted for patients consid ered 'high risk' and 'low risk' for recurrence based upon the following prognostic factor profiles. High risk women were as sumed to be aged 40, with four positive lymph nodes, and an ER negative grade 3 tumour 4 cm in diameter. On the basis of these factors and assuming radiotherapy to be given, the baseline prognosis for this group (i.e. the predicted probability of remaining recurrence free at 10 years in the absence of chemotherapy (and, given the negative ER status of the group, anti hormone therapy)), was estimated to be 18%. Low risk women were assumed to be aged 40, to be axillary lymph node negative, and with an ER positive grade 1 tumour 2 cm in diameter. On the basis of these factors and again assuming radiotherapy to be given, baseline prognosis (i.e. the predicted probability of remaining recurrence free at 10 years in the ab sence of chemotherapy (but, given the positive ER status of the group, in the presence of 5 years of tamoxifen)), was esti mated to be 85%. Analyses exploring changes to the age of pa tients in these high and low risk groups were also conducted.

Given the impracticalities of reporting detailed results for large numbers of differing prognostic groups, the sub group analyses provide an insight into how the cost effectiveness of chemotherapy differs between patient groups at opposite ends of the risk spectrum. ICERs from the model were,

Table 3 – Expected discounted lifetime costs (2009 £ UK) and QALYs for each of the four treatment strategies modelled based on the reference case cohort.^a

	No chemotherapy	CMF chemotherapy	E CMF/FEC60 chemotherapy	FEC D chemotherapy
Costs				
Chemotherapy ^b		£3,113	£3,691	£7,111
Routine follow up ^c	£1,087	£1,220	£1,245	£1,264
Recurrence	£13,116	£10,743	£10,310	£9,952
Total costs	£14,204	£15,076	£15,246	£18,327
Total QALYs	10.93	12.35	12.66	12.88

^a Patient is age 40, with one positive node, and a grade 2, ER negative tumour 3 cm in diameter.

^b Includes chemotherapy drug and toxicity costs.

^c Costs incurred in disease free health state.

however, also plotted against predicted baseline risk to illus trate how the most cost effective strategy changes as a func tion of the risk of recurrence.

2.7. Sensitivity analysis

Model parameters were assigned probability distributions to describe the range of plausible values they could take based on the precision of the source evidence (for further details see web Appendix 1). Probabilistic sensitivity analysis (PSA) was then conducted to examine the joint impact of all uncer tain parameters in the model upon cost effectiveness re sults.^{26–28} By running the model 3000 times, each time randomly sampling a set of plausible values from the

parameter distributions and recalculating the results, it was possible to describe the strength of evidence in favour of each treatment strategy being most cost effective. Results are presented by describing, for a range of different cost effectiveness thresholds, the probability that a strategy is cost effective.

One way sensitivity analysis was used to assess the im pact on cost effectiveness results of changes in individual model parameters and assumptions. Amongst the analyses conducted (see column 1 of Table 4 for details) were assess ments of the sensitivity of the model's results to the use of time dependent chemotherapy treatment effects, alternative durations of chemotherapy treatment effect, and an assump tion that HRQoL returns to general population levels in

Table 4 – Scenarios modelled in sub-group / one-way sensitivity analyses. Shown are associated ICERs for each treatment strategy compared with the next best alternative (the first strategy in the column to the left which is not dominated)

	ICER (Probability that strategy is cost effective at £20,000 per QALY ^a)			
	No chemotherapy	CMF chemotherapy	E CMF/FEC60 chemotherapy	FEC D chemotherapy
Reference case results		, C`		
Average risk woman aged 40 years and ER negative ^b	(0)	Dom (0)	£603 (0.28)	£13,704 (0.72)
Sub group analyses		Chi.		,
Average risk altering age and ER status				
Average risk woman aged 60 years and ER negative ^b	(0)	Dom (0)	£4,172 (0.46)	£18,550 (0.54)
Average risk woman aged 40 years and ER positive ^b	(0)	Dom (0)	£1,730 (0.66)	£24,107 (0.34)
Average risk woman aged 60 years and ER positive ^b	(0.23)	Dom (0)	£14,324 (0.74)	£45,918 (0.03)
High risk altering age		.,		
High risk woman aged 40 years and ER negative	(0)	Dom (0)	£249 (0.12)	£8,770 (0.88)
High risk woman aged 60 years and ER negative	()) </td <td>Dom (0)</td> <td>£2,317 (0.21)</td> <td>£11,195 (0.79)</td>	Dom (0)	£2,317 (0.21)	£11,195 (0.79)
Low risk altering age				
Low risk woman aged 40 years and ER positive ^d	< (0.01)	Dom (0.02)	£7,151 (0.97)	£70,116 (0)
Low risk woman aged 60 years and ER positive	(1)	Dom (0)	Dom (0)	£539,470 (0)
One way sensitivity analyses ^e			.,	
Treatment effects				
Chemotherapy treatment effects are time dependent ^f	(0)	Dom (0)	£1,219 (0.23)	£12,128 (0.77)
All chemotherapy treatment effects	(0.36)	Dom (0.06)	£17,383 (0.58)	£163,464 (0)
maintained for 2 years				
All chemotherapy treatment effects	(0)	Dom (0)	£2,031 (0.72)	£25,785 (0.28)
maintained for 5 years				
All chemotherapy treatment effects	Dom (0)	Dom (0)	(0.13)	£7,943 (0.87)
maintained for lifetime				
Treatment effect for FEC D estimated	(0)	Dom (0)	£603 (0.66)	£31,220 (0.34)
from TACT only ^g				
Toxicity related mortality				
No chemotherapy related deaths	(0)	Dom (0)	£615 (0.28)	£13,616 (0.72)
High risk of chemotherapy related death (2%)	(0)	£530 (0)	£580 (0.28)	£14,094 (0.72)
Health related quality of life				
Decrement with chemotherapy equal for all regimens	(0)	Dom (0)	£588 (0.24)	£12,996 (0.76)
(0.057) and does not continue beyond treatment				
HRQoL returns to age adjusted population norm	(0)	Dom (0)	£624 (0.29)	£14,178 (0.71)
levels for women surviving 15 years post recurrence				

Dom strategy is dominated, that is it has lower QALYs and greater costs than another strategy or a combination of other strategies.

^a Cost effectiveness acceptability curves showing results for a wider range of cost effectiveness threshold values can be found in web Appendix 1.

^b Average risk women have one positive lymph node and grade 2 tumours 3 cm in size.

^c High risk women have four positive lymph nodes, and grade 3 tumours 4 cm in size.

^d Low risk women are axillary lymph node negative with grade 1 tumours 2 cm in size.

^e Shown for the reference case cohort.

^f Analysis utilises the following time dependent annual recurrence rate ratios for CMF chemotherapy in women <50 years of age: years 0 1 = 0.52, (SE = 0.05), years 2 4 = 0.69 (SE = 0.06), years 5 9 = 0.79 (SE = 0.08).¹

^g Hazard ratio for DFS with FEC D from the TACT trial is 0.95 (95% CI: 0.85 1.08).³

women surviving 15 years post recurrence. The impact of changes to the magnitude of the treatment effect for FEC D was also explored. In this analysis, the base case treatment effect estimate for FEC D (Table 2) was replaced with the treatment effect estimate from TACT (HR for disease free sur vival 0.95 (SE 0.06)), the largest and most recent trial of sequential docetaxel and one undertaken in the UK.³

Finally, and because the European patent for docetaxel ex pired in November 2010, threshold analysis was conducted to determine the price to which the drug would need to fall in order for FEC D to be the most cost effective treatment strat egy (assuming a threshold of £20,000 per QALY) for all patient sub groups modelled.²⁹ Further sensitivity analyses are re ported in web Appendix 1.

Results

3.1. Reference case cohort

For the reference case cohort (for whom the predicted proba bility of remaining recurrence free out to 10 years in the ab sence of chemotherapy and (given the ER negative status of the group) anti hormone therapy is 44%), expected costs and QALYs estimated for the four treatment strategies are shown in Table 3 and Fig. 3. FEC D on average generates the greatest amount of health gain but also has the greatest cost. When compared with E CMF/FEC60 (the next most effective alternative) the expected lifetime incremental costs and QALYs with FEC D are £3,081 and 0.225 respectively, giving an ICER of £13,704. This value is less than the standard threshold employed by NICE (of between £20,000 and £30,000 per QALY) indicating that on average, for this patient group, FEC D is the most cost effective treatment strategy.

If the threshold is £20,000 per QALY then the probability that FEC D is cost effective is 0.72, rising to 0.85 for a thresh old of £30,000 per QALY.

3.2. Sub-group analysis

Table 4 shows the ICER for each strategy compared with its next best alternative and the probability that each strategy is cost effective at a threshold of £20,000 per QALY (results for other threshold values are shown in web Appendix 1), for the various sub groups considered. Given below in paren theses alongside the results for each sub group examined is the predicted probability of remaining recurrence free out to 10 years (recurrence free survival RFS) in the absence of che motherapy and the presence of other routinely administered adjuvant therapies (i.e. radiotherapy and anti hormone ther apy for ER positive patients). Taking the reference cohort first, FEC D remained cost effective when patient age was in creased from 40 to 60 years (10 year RFS of 50%). Switching ER status from negative to positive increased the ICER for FEC D versus E-CMF/FEC60 in younger women to £24,107, sug gesting that a decision about the most cost effective regimen would be finely balanced (10 year RFS of 65%). For ER positive older women the ICER for FEC D exceeded £30,000 per QALY and E CMF/FEC60 is cost effective at conventional thresholds (10 year RFS of 77%).

For women considered high risk for recurrence FEC D was the optimal treatment strategy and this remained the case when patient age was increased from 40 (10 year RFS of



	No chemotherapy	CMF	E-CMF/ FEC60	FEC-D
Incremental cost-effectiveness ratio		Dominated	£603	£13,704

Fig. 3 – Expected discounted lifetime costs and QALYs and cost-effectiveness for the reference-case cohort (age 40, 1 positive node, with a 3 cm, grade 2, ER negative tumour – predicted probability of remaining recurrence free at 10-years in absence of chemotherapy and anti-hormone therapy of 44%).

18%) to 60 years (10 year RFS of 22%). For younger low risk wo men (10 year RFS of 85%) the ICER for FEC D exceeded £70,000 per QALY, and E CMF/FEC60 was the most cost effective strat egy. For older low risk women (10 year RFS of 92%) a policy of no chemotherapy was most cost effective, proving to be less costly and associated with better quality of life than both CMF and E CMF/FEC60. The ICER for FEC D versus no chemo therapy in this group exceeded £539,000 per QALY.

Fig. 4a and b further explore, for ER negative and positive patients respectively, the potential for cost effectiveness to

vary by baseline prognosis and age. In both figures, the ICER for FEC D versus E CMF/FEC60 increases (FEC D becomes less cost effective) as baseline prognosis improves (baseline prog nosis is the probability of remaining recurrence free out to 10 years in the absence of chemotherapy, as predicted by the model on the basis of established prognostic factors plus other routinely administered adjuvant therapies (i.e. radio therapy, and anti hormone therapy for ER positive women)). Assuming a cost effectiveness threshold of £20,000 per QALY, Fig. 4a and b can be used to infer the baseline prognosis



ER positive aged 40 years ER positive aged 60 years

Fig. 4 – Incremental cost-effectiveness ratio (ICER) for FEC-D versus E-CMF plotted against baseline prognosis and by age group, for ER negative (a) and ER positive (b) patients'. * ER positive women aged 40 years are assumed to receive five years of anti-hormone therapy with tamoxifen. ER positive women aged 60 years are assumed to receive five years of anti-hormone therapy with an aromatase inhibitor.

beyond which FEC D is unlikely to be cost effective (i.e. the ICER exceeds the £20,000 threshold). The figures show that within each age group, the baseline prognoses at which the most cost effective treatment switches from FEC D to E CMF/FEC60 is similar for ER negative and ER positive pa tients. For 40 year old ER negative women, for example (Fig. 4a), FEC D is likely to be cost effective until baseline prognosis exceeds 59%. For 40 year old ER positive women (Fig. 4b), baseline prognosis must exceed 60% before FEC D is unlikely to be cost effective. For older women, in whom chemotherapy is less effective, FEC D is likely to be cost effective until baseline prognosis exceeds 53% in ER negative patients (Fig. 4a) and 54% in ER positive patients (Fig. 4b). Also indicated in Fig. 4a and b are the 10 year probabilities of remaining recurrence free in the absence of chemotherapy, at which E CMF/FEC60 is replaced by 'No chemotherapy' as the optimal treatment strategy for older women. For no pa tient subgroup was CMF ever the cost effective alternative.

3.3. Sensitivity analysis

Table 4 shows that, for the reference cohort, using time dependent chemotherapy treatment effects had little impact upon the results. Other analyses showed FEC D would not be cost effective if chemotherapy reduced the risk of recur rence for just the first 2 years following surgery (the effect ceasing thereafter) (ICER £163,464), but would be highly cost effective if the treatment effect was maintained over a patient's lifetime (ICER £7,943). Running the model using the hazard ratio for FEC D versus E CMF/FEC60 for disease free survival, reported by the TACT trial, increased the base case ICER for FEC D to £31,220.

Threshold analysis showed that, for ER positive women aged 40 years and with an average risk of recurrence, the price of docetaxel would need to fall from £1232 per cycle to £1075 per cycle in order for FEC D to replace E CMF/FEC60 as the optimal treatment strategy. For ER positive women aged 60 years the price would need to fall to £700 per cycle. For low risk ER positive women aged 40 years, the price fall re quired was even greater, from £1232 to £540 per cycle. FEC D was never cost effective for low risk ER positive women aged 60 years, even when the price of docetaxel was reduced to £0.

4. Discussion

This study used decision analytic modelling to estimate and compare the lifetime costs and effects of no chemotherapy and first, second, and third generation adjuvant chemother apy for women with early breast cancer. Sub group analyses showed different treatment strategies have the potential to be cost effective for different types of patients. Assuming the cost effectiveness threshold to be £20,000 per QALY, and summarising for younger women first, FEC D appears cost effective for patients, who, on the basis of their prognostic factors and other adjuvant therapies administered (radiother apy, and in the case of ER positive women, tamoxifen) have a 10 year predicted probability of remaining recurrence free in the absence of chemotherapy of less than 59 60%. Above this range E CMF/FEC60 appears to be the cost effective alternative. Amongst older women, FEC D appears cost effec tive for patients, who, on the basis of their prognostic factors and other adjuvant therapies (radiotherapy, and in the case of ER positive women, an aromatase inhibitor) have a 10 year predicted probability of remaining recurrence free in the ab sence of chemotherapy of less than 53 54%. Above this range, E CMF/FEC60 is the cost effective alternative up until pre dicted 10 year recurrence free survival exceeds around 80% (Fig. 4a and b), at which point a policy of no chemotherapy be comes optimal.

One way sensitivity analyses demonstrated the model's results to be robust to changes in the risk of chemotherapy related mortality and the impact of adjuvant therapy on HRQoL. Results were sensitive, however, to the duration and magnitude of chemotherapy treatment effects. In particular, replacing the pooled treatment effect estimate for FEC D ver sus E CMF/FEC60 from the TACT and PACS 01 trials with the treatment effect estimate from the TACT trial alone (the larg est study and one undertaken in the UK), increased the ICER for FEC D in the reference case cohort from £13,704 to £31,220. As the newest of the three regimens compared here, the evidence base for FEC D is limited. Furthermore the two RCTs contributing effectiveness data to this study reported variable findings PACS01 finding in favour of FEC D (HR for DFS 0.80, 95% CI 0.67 0.96), TACT showing less favourable results (HR for DFS 0.95, 95% CI 0.85 1.08).^{2,3} Further consid eration of the evidence on the effectiveness of FEC D is clearly required and, until these data become available, the results reported in this paper should be interpreted with caution.

In anticipation that the cost of docetaxel will fall following Its patent expiration, threshold analysis was used to identify the cost of the drug at which FEC D would become cost effec tive for various sub groups shown in Table 4. For some groups the cost reduction required was small (for example £157 per cycle for young women with ER positive tumours and an aver age risk of recurrence). For others (older women with ER posi tive tumours and a low risk of recurrence) the model suggested that FEC D was unlikely to be cost effective, even if the cost of docetaxel was zero. This was because the incre mental cost of FEC D over no chemotherapy (attributable to docetaxel related adverse events) remained substantial in relation to the small QALY gain received by this low risk group.

This study has a number of strengths. The 'prognostic' model used to predict the risk of a first recurrent event in the absence of anti hormone therapy and chemotherapy showed a high degree of consistency with the results of the ABC and TACT trials (see web Appendix 1). Despite its wide spread use by clinicians in the UK, Adjuvant! Online was not used to generate the prognosis predictions for the health economic model.³⁰ Adjuvant!, a web based prognosis predic tion algorithm developed in the United States, provides infor mation on the 10 year cumulative risk of recurrence, rather than annual risks over a lifetime horizon as required by the disease modelling here. Furthermore, recent evidence is avail able to suggest that Adjuvant! Online may over estimate prognosis (overall and breast cancer specific survival) for wo men with early breast cancer in the UK.³¹

In populating the model, it was possible to draw upon indi vidual patient data captured during the ABC, NEAT and TACT

RCTs. All three trials elicited information on HRQoL from pa tients during and immediately following chemotherapy; the EQ 5D data collected during ABC and TACT have not previ ously been published. Such data are traditionally scarce, as illustrated by the need for many economic models of adju vant therapy to rely upon expert clinical opinion or simple assumption as the source of their HRQoL estimates.^{32–36} Here, analyses of trial data not only provided estimates of the decrements in HRQoL experienced during chemotherapy treatments, but also indicated that the negative effect of treatments continued for at least a further year. This is a no vel finding which, to our knowledge, has not been incorpo rated in any previous economic analysis in this field.

The study is, of course, not without its limitations. The chemotherapy regimens evaluated were those specifically investigated by three large UK led RCTs and so may not nec essarily reflect routine practice elsewhere. This is particularly likely to be the case with FEC D, a regimen which is not cur rently licensed for use in the UK. Decision makers will need to determine whether the results reported here are likely to be generalisable to their own settings.

One must also acknowledge that whilst breast cancer is a complex and evolving disease, decision models are essen tially simplifications. Assumptions inevitably have to be made and inputs based upon data available at the time of analysis. With evidence of improving prognoses over the last 3 decades and with the longer term follow up of patients likely to generate new information on the ways in which the disease and indeed adjuvant chemotherapy can affect dif ferent types of women, implications for the work presented here must be considered. Firstly, and in relation to improved prognoses, it could be argued that predictions of the risk of recurrence within our study may now be unrepresentative, as the Oxford Churchill Hospital dataset which informed this modelling contained patients treated between 1986 and 2001. Given that the model was able to accurately predict recur rence free survival for patients in both the ABC and TACT tri als (see web Appendix 1), we would suggest this is not the case. However, for readers for whom this is a concern, Fig. 4a and b provide a facility for reading off cost effective ness results for lower levels of risk. Secondly, it is possible that the effects of chemotherapy are modified by factors in addition to patient age (for example by ER status); as such evi dence becomes available, the model and its results could eas ily be amended to incorporate these further treatment interactions.

Although the ABC, NEAT, and TACT trials were largely con ducted before the widespread use of adjuvant anti HER 2 therapy, it is necessary to consider the potential implications of the use of HER 2 expression as a prognostic factor and pre diction of response to anti Her 2 therapy. Simplistically, in the adjuvant setting the use of trastuzumab neutralises the ad verse prognostic effect of HER 2 overexpression. Time will tell if the promising reduction in recurrence with adjuvant trast uzmab in early analyses translate into long term survival ben efits of a similar magnitude. As we assume that all HER 2 overexpressing tumours will now receive appropriate adju vant anti HER 2 therapy we have not included HER 2 expression in the model. For metastatic disease, the cost of anti HER 2 treatment (in combination with paclitaxel) can exceed £25,000 per annum, thereby adding substantially to treatment costs post recurrence.³⁷ By preventing more of these recurrences, and thereby the need for health care providers to purchase anti Her 2 therapy to treat recurrent disease, adjuvant chemotherapy may appear more cost effective than shown here. Further research is warranted to determine the net impact of trastuzumab for advanced breast cancer, on the results presented here.

5. Conclusion

This paper has presented the first known comparison of the cost effectiveness of three different generations of adjuvant chemotherapy for early breast cancer patients with differing prognoses. The results are necessarily complex, with cost effectiveness shown to vary not only with the underlying risk of a recurrence, but also according to age and ER status. The findings are also subject to uncertainty, particularly in rela tion to the effectiveness of FEC D, a parameter on which fur ther research is clearly required.

At a time when health care resources are particularly scarce and there is a need to identify and target increasingly expensive therapies towards those patients for whom the benefits justify the costs, our findings, based upon best avail able data, should prove useful for decision makers formulat ing cost effective adjuvant chemotherapy treatment protocols in the field of early breast cancer.

Funding

Cancer Research UK (C14185/A5946), UK Medical Research Council.

Individual contributions

HC developed, populated, and analysed the economic model and drafted the manuscript. DE assisted in the development of the economic model, conducted analyses of trial data for use in the model, and helped draft the manuscript. DB was the clinical lead for the project, provided input into the mod elling process, helped write the manuscript, and co ordinated manuscript circulation across the authorship. SG advised on the development and population of the economic model and provided comments on multiple versions of the manu script. AM provided technical advice on economic model development and population and provided feedback on the manuscript. MS secured funding for the study, co ordinated the work, advised on all aspects of the modelling, and com mented upon all drafts of the manuscript produced. AG and AH supervised HC during the development of the prognostic model, and provided comments on the manuscript. JB, HE, PH, LJ, PBL, DC, JY, CP, LH, JD, and PE, provided comments on the manuscript and were pivotal in the design and conduct of the clinical trials on which the analysis was based.

Conflict of interest statement

HC had access to all data used in this study and had final responsibility for the decision to submit the paper for

publication. DB received educational travel grants from Roche and Astra Zeneca. DC received research funding, honoraria, consultancies and travel grants from Pfizer and Sanofi Aventis. PBL in the past received an honorarium for roles on Advisory boards of Sanofi Aventis and Pfizer. HE has received unrestricted educational grants for research trials from Pfizer (epirubicin), and Sanofi Aventis (docetaxel). MS is a director and minority shareholder of a consultancy company which undertakes work for pharmaceutical companies. He has not personally been involved in consultancy projects relating to the chemotherapy regimens evaluated in this study.

Role of funding source

The funding source had no involvement in the study design or analysis, the interpretation of the results, the writing of the paper, or the decision to submit for publication.

Acknowledgements

We would like to thank all of the staff who worked on the ABC, NEAT, and TACT trials, and in the Oxford Breast Care team, including surgeons, oncologists, pathologists, nurses, data managers, QoL study coordinators, support staff, and N Atkins (ICR CTSU) for helping to prepare the manuscript for submission. We are also grateful to the two anonymous reviewers for their helpful and constructive comments. Above all though, we would like to thank the patients themselves. The research costs of the ABC trial were met by Cancer Re search UK and the UK Medical Research Council (MRC), for the NEAT trial, by Cancer Research UK and Pfizer, and for the TACT trial, by Cancer Research UK and through educa tional grants from Sanofi Aventis, Pfizer and Roche. Cancer Research UK funded data management and the Oxford NHS Comprehensive Biomedical Research Centre funded data fol low up, at the Churchill Hospital. The health economic mod elling work was funded by Cancer Research UK and by the MRC through a special training fellowship in Health Services Research held by Helen Campbell.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ejca.2011.06.019.

REFERENCES

- 1. Earyl Breast Cancer Trialists' Collaborative Group. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15 year survival: an overview of the randomised trials. *Lancet* 2005;**365**(9472):1687 717.
- Roche H, Fumoleau P, Spielmann M, et al. Sequential adjuvant epirubicin based and docetaxel chemotherapy for node positive breast cancer patients: the FNCLCC PACS 01 Trial. J Clin Oncol 2006;24(36):5664 71.
- 3. Ellis P, Barrett Lee P, Johnson L, et al. Sequential docetaxel as adjuvant chemotherapy for early breast cancer (TACT): an

open label, phase III, randomised controlled trial. *Lancet* 2009;**373**(9676):1681 92.

- 4. Poole CJ, Earl HM, Hiller L, et al. Epirubicin and cyclophosphamide, methotrexate, and fluorouracil as adjuvant therapy for early breast cancer. N Engl J Med 2006;**355**(18):1851 62.
- 5. The Adjuvant Breast Cancer Trials Collaborative Group. Polychemotherapy for early breast cancer: results from the international adjuvant breast cancer chemotherapy randomized trial. J Nat Cancer Inst 2007;**99**(7):506 15.
- 6. Ward S, Simpson E, Davis S, et al. Taxanes for the adjuvant treatment of early breast cancer: systematic review and economic evaluation. *Health Technol Assess* 2007. Report No.: 11(40).
- 7. Bristol Myers Squibb. Submission to the National Institute for Health and Clinical Excellence. Taxol (Paclitaxel) for the adjuvant treatment of early breast cancer, London: National Institute for Health and Clinical Excellence; 2006.
- Sanofi Aventis. Docetaxel for the adjuvant treatment of early node positive breast cancer. Single technology appraisal to the National Institute for Health and Clinical Excellence. London: National Institute for Health and Clinical Excellence; 2006.
- Marino P, Siani C, Roche H, et al. Cost effectiveness of adjuvant docetaxel for node positive breast cancer patients: results of the PACS 01 economic study. Ann Oncol 2010;21(7):1448 54.
- Ferguson T, Wilcken N, Vagg R, et al. Taxanes for adjuvant treatment of early breast cancer. Cochrane Database of Syst Rev 2007; (Issue 4):Art. No.: CD004421.
- 11. Chang JC, Hilsenbeck SG. Prognostic and predictive markers. In: Harris JR, Lippman ME, Morrow M, et al. editors. Diseases of the breast. 3rd ed. Philadelphia: Lippincott, Williams and Wilkins; 2004. p. 675 96.
- 12. Blamey RW, Davies CJ, Elston CW, et al. Prognostic factors in breast cancer the formation of a prognostic index. Clin Oncol 1979;5(3):227 36.
- 13. Walker R. Prognostic and predictive factors in breast cancer. 1st ed. London: Martin Dunitz; 2003.
- National Institute for Health and Clinical Excellence. Docetaxel for the adjuvant treatment of early node positive breast cancer. NICE technology appraisal guidance 109. London: National Institute for Health and Clinical Excellence; 2006.
- National Institute for Health and Clinical Excellence (NICE). Guide to the Methods of Technology Appraisal. London: NICE; 2008.
- 16. Treasury HM. The green book: appraisal and evaluation in central government. London: Great Britain HM Treasury; 2003.
- 17. Euroqol Group. Euroqol a new facility for the measurement of health related quality of life. *Health Policy* 1990;**16**:199 208.
- Dolan P, Gudex C, Kind P, Williams A. A social tariff for EuroQol: results from a UK general population. Discussion paper 138. York: Centre for Health Economics, University of York; 1995.
- Campbell HE, Gray AM, Harris AH, et al. Estimation and external validation of a new prognostic model for predicting recurrence free survival for early breast cancer patients in the UK. Br J Cancer 2010;103:776 86.
- Louwman WJ, Klokman WJ, Coebergh JW. Excess mortality from breast cancer 20 years after diagnosis when life expectancy is normal. Br J Cancer 2001;84(5):700 3.
- Government Actuary's Department. Interim life tables for the United Kingdom, females based on the years 2002 2004. London: Government Actuary's Department; 2005.
- 22. Office for National Statistics. Mortality Statistics. Cause. Review of the Registrar General on deaths by cause, sex and age, in England and Wales, 2005. Series DH2 No. 32. London: Office for National Statistics; 2006.

- 23. National Institute for Clinical Excellence. Early and locally advanced breast cancer. Diagnosis and Treatment. NICE clinical guideline 80. London: NICE; 2009.
- Karnon J, Kerr GR, Jack W, et al. Health care costs for the treatment of breast cancer recurrent events: estimates from a UK based patient level analysis. Br J Cancer 2007;97:479 85.
- 25. Karlsson G, Johannesson M. The decision rules of cost effectiveness analysis. *PharmacoEconomics* 1996;**9**(2):113 20.
- 26. Briggs A, Sculpher M, Claxton K. Decision modelling for health economic evaluation. Oxford: Oxford University Press; 2006.
- Briggs AH, Goeree R, Blackhouse G, O'Brien BJ. Probabilistic analysis of cost effectiveness models: choosing between treatment strategies for gastroesophageal reflux disease. Med Decis Making 2002;22(4):290 308.
- Claxton K, Sculpher M, McCabe C, et al. Probabilistic sensitivity analysis for NICE technology assessment: not an optional extra. *Health Econ* 2005;14:339 47.
- Intellectual Property Office. Supplementary Protection Certificate Search docetaxel. http://www.ipo.gov.uk/p find spc byspc results.htm?number=SPC/GB96/009> [Accessed 12.12.10.
- Ravdin P, Siminoff L, Davis G, et al. Computer program to assist in making decisions about adjuvant therapy for women with early breast cancer. J Clin Oncol 2001;19(4):980 91.
- Campbell HE, Taylor MA, Harris AL, et al. An investigation into the performance of the ADJUVANT! Online prognostic programme in early breast cancer for a cohort of patients in the UK. Br J Cancer 2009;101(7):1074 84.
- Hillner BE, Smith TJ. Efficacy and cost effectiveness of adjuvant chemotherapy in women with node negative breast cancer: a decision analytic model. N Engl J Med 1991;324:160 8.
- Hillner BE, Smith TJ. Should women with node negative breast cancer receive adjuvant chemotherapy? Insights from a decision analysis model. Breast Cancer Res Treat 1992;23:17 27.
- Hillner BE, Smith TJ. A model of chemotherapy in node negative breast cancer. J Nat Cancer Inst Monogr 1992;11:143 9.
- Naeim A, Keeler EB. Is adjuvant therapy for older patients with node() early breast cancer cost effective. Crit Rev Oncol Hematol 2005;53:81 9.
- Naeim A, Keeler EB. Is adjuvant therapy for older patients with node(+) early breast cancercost effective? Breast Cancer Res Treat 2005;94:95 103.
- 37. National Institute for Clinical Excellence (NICE). Guidance on the use of trastuzumab for the treatment of advanced breast cancer. London: NICE; 2002.
- Caldwell D. Introduction to indirect and mixed treatment comparisons (MTC). Multiple treatment comparisons workshop for NICE. 1st September 2006. http:// www.bris.ac.uk/cobm/docs/mtc indirect.pdf>. [Accessed 29.05.10].
- Joint Formulary Committee. British National Formulary. No. 58. London: British Medical Association and Royal Pharmaceutical Society of Great Britain; 2009.
- Lober J, Sogaard J, Mouridsen HT, et al. Treatment costs of adjuvant cytotoxic therapy in premenopausal breast cancer patients. Acta Oncol 1988;27(6A):767 71.

- Lokich JJ, Moore CL, Anderson NR. Comparison of costs for infusion versus bolus chemotherapy administration: analysis of five standard chemotherapy regimens in three common tumors Part one. Model projections for cost based on charges. *Cancer* 1996;78(2):294 9.
- 42. Curtis L. Unit costs of health and social care 2009. Canterbury: The University of Kent; 2009.
 43. Department of Health NUC Reference Costs 2009.
- 43. Department of Health. NHS Reference Costs 2008 2009. London: Department of Health; 2009.
- 44. Department of Health, Joint Surveys Unit of Social and Community Planning Research, Department of Epidemiology and Public Health University College London. Health Survey for England. London: Department of Health; 1998.
- 45. Howell A, Cuzick J, Baum M, et al. Results of the ATAC (Arimidex, Tamoxifen, Alone or in Combination) trial after completion of 5 years' adjuvant treatment for breast cancer. *Lancet* 2005;**365**(9453):60 2.
- Department of Health. NHS Reference Costs 2005 06. London: Department of Health; 2006.
- 47. Suh WW, Hillner BE, Pierce LJ, Hayman JA. Cost effectiveness of radiation therapy following conservative surgery for ductal carcinoma in situ of the breast. Int J Radiat Oncol Biol Phys 2005;**61**(4):1054 61.
- Stalmeier PF. Discrepancies between chained and classic utilities induced by anchoring with occasional adjustments. Med Decis Making 2002;22(1):53 64.
- 49. Hayman JA, Fairclough DL, Harris JR, et al. Patient preferences concerning the trade off between the risks and benefits of routine radiation therapy after conservative surgery for early stage breast cancer. J Clin Oncol 1997;15:1252 60.
- 50 Sorensen S, Brown R, Benedict A, et al. Patient rated utilities in postmenopausal early breast cancer: a cross country comparison. Hamburg: ISPOR Seventh Annual European Congress; 2004.
- Cykert S, Phifer N, Hansen C. Tamoxifen for breast cancer prevention: a framework for clinical decisions. Obstet Gynecol 2004;104(3):433 42.
- Chie WC, Huang CS, Chen JH, et al. Utility assessment for different clinical phases of breast cancer in Taiwan. J Formos Med Assoc 2000;99(9):677 83.
- Grann VR, Panageas KS, Whang W, et al. Decision analysis of prophylactic mastectomy and oophorectomy in BRCA1 positive or BRCA2 positive patients. J Clin Oncol 1998;16:979 85.
- Grann VR, Jacobson JS, Sundararajan V, et al. The quality of life associated with prophylactic treatments for women with BRCA1/2 mutations. *Cancer J Sci Am* 1999;5:283 92.
- 55. Brown RE, Hutton J, Burrell A. Cost effectiveness of treatment options in advanced breast cancer in the UK. *PharmacoEconomics* 2001;19(11):1091 1102: 1102.
- 56. Hutton J, Brown RE, Borowitz M, et al. A new decision model for cost utility comparisons of chemotherapy in recurrent metastatic breast cancer. *PharmacoEconomics* 1996;9(Suppl 2):8 22.
- 57. Brown RE, Hutton J. Cost utility model comparing docetaxel and paclitaxel in advanced breast cancer patients. *Anticancer Drugs* 1998;9:899 907.