# National Communicable Diseases Surveillance Report

## Fortnight 18, 2021 Summary Notes for Selected Diseases

# 30 August to 12 September 2021

## Infectious and congenital syphilis

Increases in infectious syphilis notifications are attributed to an on-going outbreak occurring in Aboriginal and Torres Strait Islander people residing in northern and central Australia, continued increases among men who have sex with men (MSM) in urban areas, and increases in women (Aboriginal and Torres Strait Islander and non-Indigenous) residing in urban areas of Australia.

## Outbreak in northern and central Australia

In January 2011, an increase of infectious syphilis notifications among Aboriginal and Torres Strait Islander people was identified in the North West region of Queensland, following a steady decline at a national level in remote communities. Subsequent increases in infectious syphilis notifications were reported in the Northern Territory in 2013, Western Australia in 2014 and South Australia in 2016, following sustained periods of low notification rates. The outbreak is of significant public health concern given the: elevated rates of infectious syphilis among women of child-bearing age, increasing the risk of congenital syphilis; and the concomitant risk of HIV transmission. For the latest information on the infectious syphilis outbreak and related national activities, refer to the <u>Department's website</u>.

#### Increases among MSM

Since 2010 increases in notifications of infectious syphilis have been reported in MSM, predominately 20-39 years of age, residing in urban areas of Australia.

Increases among women (Aboriginal and Torres Strait Islander and non-Indigenous) Since 2016, increases in notifications of infectious syphilis have been reported in women (Aboriginal and Torres Strait Islander and non-Indigenous) aged predominately 20-39 years of age residing in urban areas in Australia. As noted in the outbreak in northern and central Australia, increases in women of childbearing age is of significant public health concern given the increased risk of congenital syphilis.

# Syphilis response

On 23 March 2021, the Australian Health Protection Principal Committee (AHPPC) endorsed the *National strategic approach for responding to rising rates of syphilis in Australia 2021* (Strategic Approach) prepared through the Communicable Diseases Network Australia (CDNA) and BBV STI Standing Committee (BBVSS). The Strategic Approach builds on and intersects with existing national activities related to syphilis and provides specific focus for efforts towards rising rates of syphilis and adverse outcomes in Australia.

The CDNA and BBVSS are, in collaboration, developing priority public health actions, including those related to workforce and community engagement, to ensure progress is made towards reducing the incidence of syphilis and elimination of congenital syphilis in Australia. These actions will be provided to AHPPC for endorsement in the coming months. For further information on national activities related to STIs, including syphilis, refer to the <u>Department's website</u>.

#### Interpretative Notes

Selected diseases are chosen each fortnight based on either exceeding two standard deviations from the 90 day and/or 365 day five year rolling mean or other disease issues of significance identified during the reporting period. All diseases reported are analysed by notification receive date. Data are extracted each Monday of a CDNA week.

Totals comprise data from all States and Territories. Cumulative figures are subject to retrospective revision so there may be discrepancies between the number of new notifications and the increment in the cumulative figure from the previous period.

<sup>1</sup>The past quarter (90 day) surveillance period includes the date range (15/06/2021 to 12/09/2021).

<sup>2</sup>The quarterly (90 day) five year rolling mean is the average of 5 intervals of 90 days up to 12/09/2021. The ratio is the notification activity in the past quarter (90 days) compared with the five year rolling mean for the same period.

<sup>3</sup>The past year (365 day) surveillance period includes the date range (13/09/2020 to 12/09/2021).

<sup>4</sup>The yearly (365 day) five year rolling mean is the average of 5 intervals of 365 days up to 12/09/2021. The ratio is the notification activity in the past year (365 days) compared with the five year rolling mean for the same period.

The five year rolling mean and the ratio of notifications compared with the five year rolling mean should be interpreted with caution. Changes in surveillance practice, diagnostic techniques and reporting may contribute to increases or decreases in the total notifications received over a five year period. Ratios are to be taken as a crude measure of current disease activity and may reflect changes in reporting rather than changes in disease activity.

AD	DT FN18/2021	State or Territory									Notification received dat Totals for Australia				Historical 90 Day Period				Historical Yearly Period			
Disease group	Disease name	Disease code	АСТ	NSW	NT	Qld	SA	Tas	Vic	WA	This reporting period 30/08/2021 12/09/2021	Previous reporting Period 16/08/2021 29/08/2021	Same reporting period last year 30/08/2020 12/09/2020	Current year YTD 01/01/2021 12/09/2021	Past Quarter 15/06/2021 12/09/2021	Quarterly rolling 5 year mean	Ratio past quarter/5 year mean*	Exceeds quarterly rolling mean +2 SD by	Past Year 13/09/2020 12/09/2021	Yearly rolling 5 year mean 13/09/2015 12/09/2020	Ratio past year/5 year mean*	Exceeds yearly rolling mean +2 S by
Bloodborne diseases	Hepatitis B (newly acquired)	039	-	-	-	1	-	· .	-	-	1	3	2	66	23	36.0	0.6	-	99	148.6	0.7	-
	Hepatitis B (unspecified) Hepatitis C (newly acquired)	052	- 4	40	-	33 18	2	- 1	48	25	153 22	173 36	176 32	3,239 529	1,039 184	1,456.0 169.0	0.7	-	4,659 714	5,809.2 709.2	0.8	-
	Hepatitis C (unspecified)	053	6	63	5	46	-	4	64	30	218	249	295	4,901	1,593	2,258.2	0.7	-	7,110	9,563.6	0.7	-
	Hepatitis D	050	-	2	-	-	-	· ·	-	1	3	2	4	57	18	19.2	0.9	-	80	68.2	1.2	-
Gastrointestinal diseases	Botulism Campylobacteriosis	045	- 18	- 236	- 11	- 272	- 117	- 34	- 276	- 99	- 1,063	- 1,176	- 1,104	25,048	1 8,072	0.2 6,940.8	5.0 1.2	-	3 36,129	1.0 30,222.2	3.0 1.2	-
	Cryptosporidiosis	061	-	3	1	14	2	1	27	3	51	63		1,371	415	486.2	0.9	-	1,744	3,821.4	0.5	-
	Haemolytic uraemic syndrome (HUS) Hepatitis A	055 038	-	- 2	-	-	-	<u> </u>	- 1	-	- 3	- 2	- 1	4 26	- 19	3.4 42.8	- 0.4	-	6 30	16.2 233.2	0.4	-
	Hepatitis E	051	-	-	-	-	-	-	-	-	-	-	1	11	4	9.2	0.4	-	12	47.2	0.1	-
	Listeriosis	018	-	-	-	-	-	•	-	-	-	2	1	30	11	13.6	0.8	-	47	67.4	0.7	-
	Paratyphoid Salmonellosis	080	- 3	- 52	- 18	- 52	- 19	- 4	- 39	- 15	- 202	- 216	- 234	3	3 1,510	9.0 2,501.0	0.3	-	3 10,422	81.8 15,417.4	0.0	-
	Shigellosis	031	-	1	4	-	4	•	1	3	13	17	27	322	93	468.0	0.2	-	534	2,099.8	0.3	-
	STEC Typhoid Fever	054	-	- 4	-	-	9	·	1	- 2	- 16	- 16	16	398 8	104 3	98.2 20.0	1.1 0.2	-	561 15	500.8 147.8	1.1 0.1	-
Quarantinable diseases	Avian influenza in humans (AIH)	035	-	-	-	-	-		-	-	-	-	-	-	-	- 20.0	0.2	-	-	- 147.0	0.1	-
	Cholera	008	-	-	-	-	-	-	-	-	-	-	-	-	-	-		-	-	1.2	-	-
	COVID-19 Middle East respiratory syndrome coronavirus (N	081 V 079	251	18,734	-	- 23	- 23	-	<u>3,743</u> -	- 2	22,776	13,295 -	1,026	47,370	45,359	3,836.8 -	11.8	24,363.5	49,198 -	<u>5,352.0</u> -	9.2	<u>19,911.1</u> -
	Plague	025	-	-	-	-	-		-	-	-	-	-	-	-	-		-	-	-		-
	Rabies	028	-	-	-	-	-	·	-	-	-	-	-	-	-	-		-	-	-		-
	Severe acute respiratory syndrome (SARS) Smallpox	0/1	-	-	-	-	-		-	-	-	-	-	-	-	-		-	-	-		-
	Viral haemorrhagic fever (NEC)	036	-	-	-	-	-	•	-	-	-	-	-	-	-	-		-	-	-		-
Sexually transmissible infections	Yellow fever Chlamydial infection	041	- 37	- 505	- 21	- 798	- 184	- 54	- 301	- 458	- 2,358	- 2,799	- 3,209	- 57,235	- 18,221	- 24,282.0	0.8	-	- 82,865	- 100,409.0	0.8	-
	Donovanosis	010	-	- 303	-	-	-	-	- 301	-	-	-	- 3,205	- 37,235	- 10,221	- 24,282.0	0.8	-	- 82,803	- 100,409.0	0.8	-
	Gonococcal infection	011	12	149	31	249	65	4	213	139	862	851	1,061	19,089	6,203	7,129.2	0.9	-	26,882	28,896.2	0.9	-
	Syphilis < 2 years Syphilis > 2 years or unspecified duration	066 067	-	- 25	6 3	38 1	- 7	-	44 27	<u>33</u> 6	<u>154</u> 37	<u>179</u> 61	215 105	3,863 1,239	1,238 362	<u>1,210.4</u> 542.8	1.0 0.7	-	5,383 1,808	4,734.0 2,145.2	1.1 0.8	-
	Syphilis congenital	047	-	-	-	-	-	•	-	-	-	1	2	11	5	2.4	2.1	-	16	7.8	2.1	-
Vaccine preventable diseases	Diphtheria	009	-	· ·	-	-	-	· ·	-	-	-	-	-	4	2	1.6	1.3	-	8	8.2	1.0	-
	Haemophilus influenzae type b Influenza (laboratory confirmed)	012	-	- 2	- 5	- 13	- 5	- 3	-	-	- 28	- 21	- 40	13 529	4	5.4 83,431.0	0.7	-	20 737	18.8 152,940.4	1.1	-
	Measles	021	-	-	-	-	-	•	-	-	-	-	-	-	-	17.4	-	-	-	121.6	-	-
	Mumps Pertussis	043	- 1	- 1	-	- 5	- 3	- 1	- 16	- 2	- 29	- 24	7 24	13 428	3 154	131.2 2,593.0	0.0	-	26 577	584.8 14,166.2	0.0	-
	Pneumococcal disease (invasive)	065	-	12	5	18	5	1	10	8	63	52	45	1,035	465	661.6	0.1	-	1,363	1,827.6	0.7	-
	Poliovirus infection	026	-	-	-	-	-	· .	-	-	-	-	-	-	-	-		-	-	-		-
	Rotavirus Rubella	077	-	12	- 13	- 20	- 14	- 1	NN -	17	82	- 60	- 34	947	390 1	1,050.8 2.0	0.4	-	1,276 3	4,541.0 12.6	0.3	-
	Rubella congenital	046	-	-	-	-	-	•	-	-	-	-	-			-		-	-	0.2	-	-
	Tetanus Varicella zoster (chickenpox)	033	- 1	- NN	- 1	- 1	- 12	-	- 12	27	1	- 59	- 113	4	1 424	0.6 975.8	1.7 0.4	-	7 2,440	4.2 3,599.4	1.7 0.7	0.2
	Varicella zoster (shingles)	074	18		9	1	110	21		81	335	347		7,409	2,196	3,039.2	0.4	-	12,048	11,913.0	1.0	-
	Varicella zoster (unspecified)	075	-	NN	6	365	45	18		70	942	878	522	13,858	5,415	3,560.8	1.5	276.5	18,270	14,067.2	1.3	170.0
Vectorborne diseases	Barmah Forest virus infection Chikungunya virus infection	048	-	- 1	-	- 8	-	· ·	-	-	- 9	- 8	- 20	283	- 74	94.0 14.6	- 0.8	-	443 2	411.6 78.4	1.1	-
	Dengue virus infection	003	-	-	-	-	-	•	-	-	-	-	1	2	-	255.6	-	-	2	1,275.0	0.0	-
	Flavivirus infection (unspecified) Japanese encephalitis virus infection	001	-	-	-	-	-	÷	-	-	-	-	-	3	-	6.2 0.4	-	-	8	32.6 1.2	0.2	-
	Malaria	039	-	- 1	-	1	-		-	-	2	4	3	34	17	80.0	0.2	-	50	340.0	0.8	-
	Murray Valley encephalitis virus infection	049	-	- 7	-	-	-	•	-	- 7	-	-	-	1	1	-		1.0	1	0.2	5.0	-
	Ross River virus infection West Nile/Kunjin virus infection	002	-	- 7	-	- 17	-	<u> </u>	2	- 7	- 36	- 44	- 64	2,847	426	592.6 0.6	0.7	-	3,619	4,699.0 1.6	0.8	-
Zoonoses	Anthrax	058	-	-	-	-	-	•	-	-	-	-	-	-	-	-		-	-	-		-
	Australian bat lyssavirus infection	063 004	-	-	-	- 2	-	-	-	-	- 2	-	-	- 16	-	- 4.8	10	-	-	-	1.1	-
	Brucellosis Leptospirosis	004	-	- 1	-	2	-	<u> </u>	-	-	3	1	- 3	16 213	5 37	4.8	1.0 1.2	-	20 238	19.0 117.0	1.1 2.0	- 53.9
	Lyssavirus infection (NEC)	064	-	-	-	-	-	· ·	-	-	-	-	-	-	-	-		-	-	-		-
	Ornithosis Q fever	023	-	- 3	-	- 7	-	-	- 1	-	1	- 16	- 13	19 362	6 90	5.2 112.8	1.2 0.8	-	52 480	23.2 536.4	2.2	9.2
	Tularaemia	027	-	-	-	-	-		-	-	-	- 10	- 15	- 502	- 90	-	0.0	-	-	0.4	-	-
Other notifiable infections	iGAS	082	-	-	-	9	-		-	2	11	12		85	75	-		75.0	85	-		85.0
	Legionellosis Leprosy	015 016	-	-	-	-	2	· ·	-	- 3	- 9	- 12	- 12	358	75 2	84.0 2.8	0.9	-	536 9	421.0 11.0	1.3 0.8	-
	Meningococcal disease (invasive)	022	-	1	-	-	-	•	1	1	3	4	1	52	14	72.6	0.2	-	76	249.2	0.3	-
	RSV	083	-	-	-	31	-	-	-	-	31	39	-	310	310	310.0		310.0	310	-		310.0
	Tuberculosis	034	1	17	-	-	-		13	4	42	46	68	993	333	1,538.0	0.9	-	1,538	1,451.2	1.1	-

Footnotes: \* Ratio of the 90 day prior surveillance period to the past 90 day 5 year rolling mean, or ratio of the year period prior surveillance period to the year period 5 year rolling mean. NN = Not Notifiable, NEC = Not Elsewhere Classified The data in this report are reliant on the provision of data from states and territories to the Australian Government Department of Health. Backlogs in notifications at the state or territory level may contribute to delays in reporting to the NNDSS. Notifications for some high volume conditions are only uploaded quarterly by some jurisdictions, which can result in apparent large variability over time. The NNDSS is a dynamic dataset, with data in this report representing data available on **(14/09/2021)**. Data in this report are subject to retrospective revision and may vary from data reported in published NNDSS reports and reports of notification data by states and territories.