



The Childhood Schedule from 1 July 2018

Birth

- Hepatitis B (usually offered in hospital)

2 months

(can be given
from 6 weeks
of age)

- Diphtheria, tetanus, whooping cough, hepatitis B, polio, *Haemophilus influenzae* type b (Hib)
- Pneumococcal
- Rotavirus

4 months

- Diphtheria, tetanus, whooping cough, hepatitis B, polio, *Haemophilus influenzae* type b (Hib)
- Pneumococcal
- Rotavirus

6 months

- Diphtheria, tetanus, whooping cough, hepatitis B, polio, *Haemophilus influenzae* type b (Hib)
- Pneumococcal—Aboriginal and Torres Strait Islander children (QLD, NT, WA, SA) and medically at-risk

12 months

- Meningococcal ACWY
- Measles, mumps, rubella
- Pneumococcal
- Hepatitis A—Aboriginal and Torres Strait Islander children (QLD, NT, WA, SA)

18 months

- Haemophilus influenzae* type b (Hib)
- Measles, mumps, rubella, chickenpox
- Diphtheria, tetanus, whooping cough
- Hepatitis A—Aboriginal and Torres Strait Islander children (QLD, NT, WA, SA)

4 years

- Diphtheria, tetanus, whooping cough, polio
- Pneumococcal—medically at-risk

- Influenza vaccine is funded for Aboriginal and Torres Strait Islander children aged 6 months to less than 5 years.
- Additional vaccines might be funded by some States and Territories.



Have questions?

Visit health.gov.au/immunisation

Make an appointment with your GP or other vaccination provider **Call 1800 671 811** for the National Immunisation Program Information Line